

VL Formal Modeling (WS 2022)


# Symbolic Summation and the modeling of sequences

Carsten Schneider

Research Institute for Symbolic Computation (RISC)  
Johannes Kepler University Linz



## General picture:



General picture:

Part 1: Symbolic summation (a short introduction)

Part 2: Modeling of sequences with a term algebra (user interface)

Part 3: Modeling of sequences in difference rings (computer algebra)

Part 4: Construction of appropriate difference rings (advanced CA)

Part 5: Applications

We start with the following telescoping problem:

**Given** an expression  $f(k)$  that evaluates to a sequence.

**Find** an expression  $g(k)$  such that the telescoping equation holds:

$$f(k) = g(k + 1) - g(k) \quad (1)$$

Suppose we find such an expression  $g(k)$ . Then we proceed as follows.

Summing (1) over  $k$  from  $a$  to  $b$  (and assuming that no poles arise during the evaluation) yields

$$\sum_{k=a}^b f(k) = g(b + 1) - g(a). \quad (2)$$

Note: we could always choose

$$g(k) = \sum_{i=a}^{k-1} f(i) \quad (3)$$

which would turn (2) to the trivial identity  $\sum_{k=a}^b f(k) = \sum_{k=a}^b f(k)$ .

Thus we should refine our problem from above:

**Find** an expression  $g(k)$  with (1) where  $g(k)$  is simpler than the trivial solution (3).

## Indefinite summation of polynomials

We start with one of the most simplest cases: the summand is a polynomial, i.e.,  $f(x) \in \mathbb{K}[x]$ .

The following questions arise:

1. What is the domain of expressions in which we search  $g(k)$ ?
2. How can we calculate a solution  $g(k)$  in this solution domain?

As it turns out, the first question can be answered nicely: a solution  $g(x)$  exists always in  $\mathbb{K}[x]$ . For the second question, we will consider two different tactics that are often used in summation packages.

## Tactic 1: the classical approach

Note that for indefinite integration of polynomials one can utilize the following well known property: for any  $m \in \mathbb{N}$  we have

$$D_x x^m = m x^{m-1}$$

which implies

$$\int_a^b x^m dx = \frac{x^{m+1}}{m+1} \Big|_a^b = \frac{b^{m+1} - a^{m+1}}{m+1}.$$

Thus by linearity we can integrate any polynomial by

$$\int_a^b \sum_{m=0}^d c_m x^m dx = \sum_{m=0}^d c_m \int_a^b x^m dx = \sum_{m=0}^d \frac{c_m (b^{m+1} - a^{m+1})}{m+1}.$$

For indefinite summation of polynomials we can follow precisely the same classical strategy.

**Definition.** For any sequence (expression)  $g(k)$  we define

$$\Delta g(k) := g(k+1) - g(k).$$

### Lemma

For  $m \in \mathbb{N}$  we have

$$\boxed{\Delta x^m = m x^{m-1}}.$$

### Proof.

We have

$$\begin{aligned}\Delta x^m &= (x+1)^m - x^m \\ &= (x+1)x(x-1)\dots(x-m+2) - x(x-1)\dots(x-m+1) \\ &= ((x+1) - (x-m+1))x(x-1)\dots(x-m+2) \\ &= m x^{m-1}.\end{aligned}$$



As a consequence we get

$$\Delta \frac{x^{m+1}}{m+1} = x^m, \quad m \in \mathbb{N}$$

and summing this equation over  $k$  from  $a$  to  $b$  yields

$$\sum_{x=a}^b x^m = \frac{(b+1)^{m+1} - a^{m+1}}{m+1}.$$

Note that this is nothing else than the continuous version for integration. In particular, for given

$$f(x) = \sum_{m=0}^d c_m x^m \in \mathbb{K}[x]$$

with  $d \in \mathbb{N}$  it follows that

$$g(x) = \sum_{m=0}^d \frac{c_m x^{m+1}}{m+1}$$

is a telescoping solution. Furthermore,

$$\sum_{x=a}^b f(x) = \sum_{m=0}^d c_m \sum_{k=a}^b k^m = \sum_{m=0}^d \frac{c_m ((b+1)^{m+1} - a^{m+1})}{m+1}.$$



The only problem is that in many cases one does not have a polynomial given in the representation of falling factorials but in the standard form

$$\sum_{m=0}^d \bar{c}_m x^m \in \mathbb{K}[x].$$

Luckily one can rewrite a polynomial written in the basis

$$1, x, x^2, \dots, x^d$$

to the representation written in the basis

$$x^{\underline{0}} = 1, x^{\underline{1}} = x, x^{\underline{2}} = x(x-1), \dots, x^{\underline{d}} = x(x-1) \dots (x-d+1)$$

by using the formula

$$x^m = \sum_{k=0}^m S(m, k) x^{\underline{k}}$$

where  $S(n, k)$  denotes the Stirling numbers of second kind. They can be computed by

$$S(n, k) = \frac{1}{k!} \sum_{i=0}^k (-1)^i \binom{k}{i} (k-i)^n;$$

**Example.** Consider the polynomial

$$f(x) = x^4.$$

Using the formulas from above, we get

$$f(x) = x^4 = \sum_{k=0}^4 S(4, k)x^k = 0x^0 + 1x^1 + 7x^2 + 6x^3 + 1x^4.$$

Thus we get

$$\begin{aligned} g(x) &= \frac{1}{2}x^2 + \frac{7}{3}x^3 + \frac{3}{2}x^4 + \frac{1}{5}x^5 \\ &= \frac{1}{30}(x-1)x(2x-1)(3x^2-3x-1). \end{aligned}$$

such that

$$g(x+1) - g(x) = f(x)$$

holds. In particular we get

$$\sum_{k=1}^n k^4 = \sum_{k=1}^n f(k) = g(n+1) - g(1) = \frac{1}{30}n(n+1)(2n+1)(3n^2+3n-1).$$

## Tactic 2: linear algebra.

We use the following property: for  $f(x) \in \mathbb{K}[x]$  there is a  $g(x) \in \mathbb{K}[x]$  with (1) where

$$\deg(g) \leq \deg(f) + 1.$$

Thus setting  $d := \deg(f) + 1$  for given  $f \in \mathbb{K}[x]$  the desired solution has the form

$$g(x) = \sum_{m=0}^d g_m x^m$$

and we can determine the unknowns  $g_0, \dots, g_d \in \mathbb{K}$  by linear algebra as follows.

**Example.** Take  $f(x) = x^4 \in \mathbb{Q}[x]$ . With  $d = \deg(f) + 1 = 5$  the ansatz

$$g(x) = g_0 + g_1x + g_2x^2 + g_3x^3 + g_4x^4 + g_5x^5$$

for the unknowns  $g_0, g_1, g_2, g_3, g_4, g_5 \in \mathbb{Q}$  is in place. This gives

$$\begin{aligned} x^4 = \Delta g(x) &= g(x+1) - g(x) = 0x^5 \\ &+ 5g_5x^4 \\ &+ (4g_4 + 10g_5)x^3 \\ &+ (3g_3 + 6g_4 + 10g_5)x^2 \\ &+ (2g_2 + 3g_3 + 4g_4 + 5g_5)x \\ &+ (g_1 + g_2 + g_3 + g_4 + g_5)x^0. \end{aligned}$$

By coefficient comparison this yields the linear system

$$\begin{array}{ll} [x^4] & 1 = 5g_5 \\ [x^3] & 0 = 4g_4 + 10g_5 \\ [x^2] & 0 = 3g_3 + 6g_4 + 10g_5 \\ [x^1] & 0 = 2g_2 + 3g_3 + 4g_4 + 5g_5 \\ [x^0] & 0 = g_1 + g_2 + g_3 + g_4 + g_5 \end{array}$$

which is already in triangular form.

Thus we can read off the solution

$$g_5 = \frac{1}{5}, \quad g_4 = -\frac{1}{2}, \quad g_3 = \frac{1}{3}, \quad g_2 = 0, \quad g_1 = -\frac{1}{30}, \quad g_0 = c$$

with  $c \in \mathbb{Q}$ . In particular, we can choose  $c = 0$  and obtain

$$g(x) = \frac{x^5}{5} - \frac{x^4}{2} + \frac{x^3}{3} - \frac{x}{30} = \frac{1}{30}(x-1)x(2x-1)(3x^2-3x-1).$$

To this end, we continue as in the previous example and get the desired result.

## More general summation objects for indefinite and definite summation

Clearly, the first tactic is very elegant, but it works only for the special case of polynomial summation. For the second tactic one has to work more (i.e., has to solve in addition a linear system), but it turns out to be more general. More precisely, one can carry over these ideas to a rather general setting that works not only for the polynomial ring  $\mathbb{Q}[x]$  but in more general rings called  $R\Pi\Sigma$ -difference rings that have been implemented within the summation package Sigma. In the following all technical details are omitted and we proceed with a concrete example.

**Example.** We want to sum

$$\sum_{k=0}^n H_k.$$

In order to accomplish this task, we take

$$f(k) = H_k$$

and search for

$$g(k) \in \mathbb{Q}(k)[H_k]$$

with

$$f(k) = g(k+1) - g(k).$$

Here we can use a similar tactic as used in the case of polynomial summation. Namely, summation theory tells us that any such solution  $g(k)$  has the property

$$\deg(g) \leq \deg(f) + 1 = 1 + 1 = 2.$$

As a consequence we can make the ansatz

$$g(k) = g_0(k)H_k^0 + g_1(k)H_k^1 + g_2(k)H_k^2$$

with  $g_0(k), g_1(k), g_2(k) \in \mathbb{Q}(k)$ .

Using recursive algorithms and linear system solving (details are skipped here) we find

$$g_0(k) = -k$$

$$g_1(k) = k$$

$$g_2(k) = 0,$$

i.e.,

$$g(k) = -k + kH_k + 0H_k^2.$$

Hence summing the telescoping equation over  $k$  from 0 to  $n$  gives

$$\sum_{k=0}^n H_k = g(n+1) - g(0) = (n+1)H_{n+1} - (n+1) = -n + (1+n)H_n.$$



The above machinery can be carried out within the summation package `Sigma`. After loading it into Mathematica

```
In[1]:= << Sigma.m
```

```
Sigma - A summation package by Carsten Schneider © RISC-JKU
```

one can insert the above sum

```
In[2]:= mySum = SigmaSum[SigmaHNumber[k], {k, 0, 1}]
```

$$\text{Out[2]} = \sum_{k=0}^n H_k$$

and can apply the command

```
In[3]:= SigmaReduce[mySum]
```

$$\text{Out[3]} = -n + (1 + n)H_n$$

In general one can insert, e.g., a sum of the form

$$\sum_{k=l}^n f(k)$$

with  $l \in \mathbb{N}$  where  $f(k)$  itself is given in terms of indefinite nested sums defined over hypergeometric products.

### Definition

Let  $\mathbb{K}$  be a field. A product  $\prod_{j=l}^k f(j)$ ,  $l \in \mathbb{N}$ , is called **hypergeometric in  $k$  over  $\mathbb{K}$**  if  $f(x) \in \mathbb{K}(x)$  is a rational function where the numerator and denominator of  $f(j)$  are nonzero for all  $j \in \mathbb{Z}$  with  $j \geq l$ . An **expression in terms indefinite of nested sums over hypergeometric products in  $k$  over  $\mathbb{K}$**  is composed recursively by the three operations  $(+, -, \cdot)$  with

- ▶ elements from the rational function field  $\mathbb{K}(k)$ ,
- ▶ hypergeometric products in  $k$  over  $\mathbb{K}$ ,
- ▶ and sums of the form  $\sum_{j=l}^k f(j)$  with  $l \in \mathbb{N}$  where  $f(j)$  is an expression in terms of indefinite nested sums over hypergeometric products in  $j$  over  $\mathbb{K}$ ; here it is assumed that the evaluation of  $f(j)$  for all  $j \geq l$  does not introduce any poles.

In[4]:= **mySum** =

**SigmaSum**[**SigmaPower**[-1, k]**SigmaBinomial**[n, k]**SigmaHNumber**[k], {k, a, b}]

$$\text{Out[4]} = \sum_{k=a}^b (-1)^k \binom{n}{k} H_k$$

In[5]:= **SigmaReduce**[**mySum**]

$$\text{Out[5]} = \left( \frac{(a-n)(-1+a-n)}{an^2} + \frac{(-1+a-n)H_a}{n} \right) (-1)^{1+a} \binom{n}{-1+a} + \left( \frac{-b+n}{n^2} + \frac{(-b+n)H_b}{n} \right) (-1)^b \binom{n}{b}$$


In[6]:= **mySum** = **SigmaSum**[**SigmaSum**[**SigmaBinomial**[n, k], {k, 0, r}]<sup>2</sup>, {r, 0, b}]

$$\text{Out[6]} = \sum_{r=0}^b \left( \sum_{k=0}^r \binom{n}{k} \right)^2$$

In[7]:= **SigmaReduce**[**mySum**]

$$\text{Out[7]} = (-b+n) \binom{n}{b} \sum_{i_1=0}^b \binom{n}{i_1} + \frac{1}{2}(2+2b-n) \left( \sum_{i_1=0}^b \binom{n}{i_1} \right)^2 - \frac{1}{2}n \sum_{i_1=0}^b \binom{n}{i_1}^2$$

## General picture:



## General picture:

Part 1: Symbolic summation (a short introduction)

Part 2: Modeling of sequences with a term algebra (user interface)

Part 3: Modeling of sequences in difference rings (computer algebra)

Part 4: Construction of appropriate difference rings (advanced CA)

Part 5: Applications

**The ground field (throughout this talk):**  $\mathbb{G} = \mathbb{K}(x)$

**The ground field (throughout this talk):**  $\mathbb{G} = \mathbb{K}(x)$

- ▶ For any element  $f = \frac{p}{q} \in \mathbb{G}$  with  $p, q \in \mathbb{K}[x]$  where  $q \neq 0$  and  $p, q$  being coprime we define

$$\text{ev}(f, k) = \begin{cases} 0 & \text{if } q(k) = 0 \\ \frac{p(k)}{q(k)} & \text{if } q(k) \neq 0. \end{cases}$$

**The ground field (throughout this talk):**  $\mathbb{G} = \mathbb{K}(x)$

- ▶ For any element  $f = \frac{p}{q} \in \mathbb{G}$  with  $p, q \in \mathbb{K}[x]$  where  $q \neq 0$  and  $p, q$  being coprime we define

$$\text{ev}(f, k) = \begin{cases} 0 & \text{if } q(k) = 0 \\ \frac{p(k)}{q(k)} & \text{if } q(k) \neq 0. \end{cases}$$

- ▶ We define  $L(f)$  to be the minimal value  $\delta \in \mathbb{N}$  such that  $q(k) \neq 0$  holds for all  $k \geq \delta$ ; further,

$$Z(f) = \max(L(\frac{1}{p}), L(\frac{1}{q})) \quad \text{if } f \neq 0.$$



**The ground field (throughout this talk):**  $\mathbb{G} = \mathbb{K}(x)$

- ▶ For any element  $f = \frac{p}{q} \in \mathbb{G}$  with  $p, q \in \mathbb{K}[x]$  where  $q \neq 0$  and  $p, q$  being coprime we define

$$\text{ev}(f, k) = \begin{cases} 0 & \text{if } q(k) = 0 \\ \frac{p(k)}{q(k)} & \text{if } q(k) \neq 0. \end{cases}$$

- ▶ We define  $L(f)$  to be the minimal value  $\delta \in \mathbb{N}$  such that  $q(k) \neq 0$  holds for all  $k \geq \delta$ ; further,

$$Z(f) = \max(L(\frac{1}{p}), L(\frac{1}{q})) \quad \text{if } f \neq 0.$$

**Example:** For

$$f = \frac{p}{q} = \frac{x - 4}{(x - 3)(x - 1)}$$

we get

$$(\text{ev}(f, n))_{n \geq 0} = (-\frac{4}{3}, \underline{0}, 2, \underline{0}, 0, \frac{1}{8}, \dots) \in \mathbb{Q}^{\mathbb{N}}$$

For  $n \geq L(f) = 4$  no poles arise;

for  $n \geq Z(f) = \max(L(\frac{1}{p}), L(\frac{1}{q})) = \max(4, 5) = 5$  no zeroes arise.

**The ground field (throughout this talk):**  $\mathbb{G} = \mathbb{K}(x)$

- ▶ For any element  $f = \frac{p}{q} \in \mathbb{G}$  with  $p, q \in \mathbb{K}[x]$  where  $q \neq 0$  and  $p, q$  being coprime we define

$$\text{ev}(f, k) = \begin{cases} 0 & \text{if } q(k) = 0 \\ \frac{p(k)}{q(k)} & \text{if } q(k) \neq 0. \end{cases}$$

- ▶ We define  $L(f)$  to be the minimal value  $\delta \in \mathbb{N}$  such that  $q(k) \neq 0$  holds for all  $k \geq \delta$ ; further,

$$Z(f) = \max(L(\frac{1}{p}), L(\frac{1}{q})) \quad \text{if } f \neq 0.$$

- ▶ We define

$$\mathcal{R} = \{r \in \mathbb{K} \setminus \{1\} \mid r \text{ is a root of unity}\}$$

with the function  $\text{ord} : \mathcal{R} \rightarrow \mathbb{Z}_{\geq 1}$  where

$$\text{ord}(r) = \min\{n \in \mathbb{Z}_{\geq 1} \mid r^n = 1\}.$$

$\mathbb{G} \longrightarrow \text{SumProd}(\mathbb{G})$  (nested sums over hypergeometric products)

Let  $\otimes$ ,  $\oplus$ ,  $\odot$ ,  $\text{Sum}$ ,  $\text{Prod}$  and  $\text{RPow}$  be operations with the signatures

$$\begin{array}{lll}
 \otimes : & \text{SumProd}(\mathbb{G}) \times \mathbb{Z} & \rightarrow \text{SumProd}(\mathbb{G}) \\
 \oplus : & \text{SumProd}(\mathbb{G}) \times \text{SumProd}(\mathbb{G}) & \rightarrow \text{SumProd}(\mathbb{G}) \\
 \odot : & \text{SumProd}(\mathbb{G}) \times \text{SumProd}(\mathbb{G}) & \rightarrow \text{SumProd}(\mathbb{G}) \\
 \text{Sum} : & \mathbb{N} \times \text{SumProd}(\mathbb{G}) & \rightarrow \text{SumProd}(\mathbb{G}) \\
 \text{Prod} : & \mathbb{N} \times \text{SumProd}(\mathbb{G}) & \rightarrow \text{SumProd}(\mathbb{G}) \\
 \text{RPow} : & \mathcal{R} & \rightarrow \text{SumProd}(\mathbb{G}).
 \end{array}$$

$\text{Prod}^*(\mathbb{G}) =$  the smallest set that contains 1 with the following properties:

1. If  $r \in \mathcal{R}$  then  $\text{RPow}(r) \in \text{Prod}^*(\mathbb{G})$ .
2. If  $f \in \mathbb{G}^*$  and  $l \in \mathbb{N}$  with  $l \geq Z(f)$  then  $\text{Prod}(l, f) \in \text{Prod}^*(\mathbb{G})$ .
3. If  $p, q \in \text{Prod}^*(\mathbb{G})$  then  $p \odot q \in \text{Prod}^*(\mathbb{G})$ .
4. If  $p \in \text{Prod}^*(\mathbb{G})$  and  $z \in \mathbb{Z} \setminus \{0\}$  then  $p^{\otimes z} \in \text{Prod}^*(\mathbb{G})$ .

$\mathbb{G} \longrightarrow \text{SumProd}(\mathbb{G})$  (nested sums over hypergeometric products)

Let  $\otimes$ ,  $\oplus$ ,  $\odot$ , **Sum**, **Prod** and **RPow** be operations with the signatures

$$\begin{array}{lll} \otimes : & \text{SumProd}(\mathbb{G}) \times \mathbb{Z} & \rightarrow \text{SumProd}(\mathbb{G}) \\ \oplus : & \text{SumProd}(\mathbb{G}) \times \text{SumProd}(\mathbb{G}) & \rightarrow \text{SumProd}(\mathbb{G}) \\ \odot : & \text{SumProd}(\mathbb{G}) \times \text{SumProd}(\mathbb{G}) & \rightarrow \text{SumProd}(\mathbb{G}) \\ \text{Sum} : & \mathbb{N} \times \text{SumProd}(\mathbb{G}) & \rightarrow \text{SumProd}(\mathbb{G}) \\ \text{Prod} : & \mathbb{N} \times \text{SumProd}(\mathbb{G}) & \rightarrow \text{SumProd}(\mathbb{G}) \\ \text{RPow} : & \mathcal{R} & \rightarrow \text{SumProd}(\mathbb{G}). \end{array}$$

**Prod\***( $\mathbb{G}$ ) = the smallest set that contains 1 with the following properties:

1. If  $r \in \mathcal{R}$  then  $\text{RPow}(r) \in \text{Prod}^*(\mathbb{G})$ .
2. If  $f \in \mathbb{G}^*$  and  $l \in \mathbb{N}$  with  $l \geq Z(f)$  then  $\text{Prod}(l, f) \in \text{Prod}^*(\mathbb{G})$ .
3. If  $p, q \in \text{Prod}^*(\mathbb{G})$  then  $p \odot q \in \text{Prod}^*(\mathbb{G})$ .
4. If  $p \in \text{Prod}^*(\mathbb{G})$  and  $z \in \mathbb{Z} \setminus \{0\}$  then  $p^{\otimes z} \in \text{Prod}^*(\mathbb{G})$ .

Furthermore, we define

$$\Pi(\mathbb{G}) = \{\text{RPow}(r) \mid r \in \mathcal{R}\} \cup \{\text{Prod}(l, f) \mid f \in \mathbb{G}, l \in \mathbb{N}\}.$$

$\mathbb{G} \longrightarrow \text{SumProd}(\mathbb{G})$  (nested sums over hypergeometric products)

Let  $\otimes$ ,  $\oplus$ ,  $\odot$ ,  $\text{Sum}$ ,  $\text{Prod}$  and  $\text{RPow}$  be operations with the signatures

$$\begin{array}{lll} \otimes : & \text{SumProd}(\mathbb{G}) \times \mathbb{Z} & \rightarrow \text{SumProd}(\mathbb{G}) \\ \oplus : & \text{SumProd}(\mathbb{G}) \times \text{SumProd}(\mathbb{G}) & \rightarrow \text{SumProd}(\mathbb{G}) \\ \odot : & \text{SumProd}(\mathbb{G}) \times \text{SumProd}(\mathbb{G}) & \rightarrow \text{SumProd}(\mathbb{G}) \\ \text{Sum} : & \mathbb{N} \times \text{SumProd}(\mathbb{G}) & \rightarrow \text{SumProd}(\mathbb{G}) \\ \text{Prod} : & \mathbb{N} \times \text{SumProd}(\mathbb{G}) & \rightarrow \text{SumProd}(\mathbb{G}) \\ \text{RPow} : & \mathcal{R} & \rightarrow \text{SumProd}(\mathbb{G}). \end{array}$$

$\text{Prod}^*(\mathbb{G}) =$  the smallest set that contains 1 with the following properties:

1. If  $r \in \mathcal{R}$  then  $\text{RPow}(r) \in \text{Prod}^*(\mathbb{G})$ .
2. If  $f \in \mathbb{G}^*$  and  $l \in \mathbb{N}$  with  $l \geq Z(f)$  then  $\text{Prod}(l, f) \in \text{Prod}^*(\mathbb{G})$ .
3. If  $p, q \in \text{Prod}^*(\mathbb{G})$  then  $p \odot q \in \text{Prod}^*(\mathbb{G})$ .
4. If  $p \in \text{Prod}^*(\mathbb{G})$  and  $z \in \mathbb{Z} \setminus \{0\}$  then  $p^{\otimes z} \in \text{Prod}^*(\mathbb{G})$ .

**Example:** In  $\mathbb{G} = \mathbb{Q}(x)$  we get

$$P = \underbrace{(\text{Prod}(1, x)^{\otimes (-2)})}_{\in \Pi(\mathbb{G})} \odot \underbrace{\text{RPow}(-1)}_{\Pi(\mathbb{G})} \in \text{Prod}^*(\mathbb{G}).$$

$\mathbb{G} \longrightarrow \text{SumProd}(\mathbb{G})$  (nested sums over hypergeometric products)

**SumProd**( $\mathbb{G}$ ) = the smallest set containing  $\mathbb{G} \cup \text{Prod}^*(\mathbb{G})$  with:

1. For all  $f, g \in \text{SumProd}(\mathbb{G})$  we have  $f \oplus g \in \text{SumProd}(\mathbb{G})$ .
2. For all  $f, g \in \text{SumProd}(\mathbb{G})$  we have  $f \odot g \in \text{SumProd}(\mathbb{G})$ .
3. For all  $f \in \text{SumProd}(\mathbb{G})$  and  $k \in \mathbb{Z}_{\geq 1}$  we have  $f^{\Delta k} \in \text{SumProd}(\mathbb{G})$ .
4. For all  $f \in \text{SumProd}(\mathbb{G})$  and  $l \in \mathbb{N}$  we have  $\text{Sum}(l, f) \in \text{SumProd}(\mathbb{G})$ .

$\mathbb{G} \longrightarrow \text{SumProd}(\mathbb{G})$  (nested sums over hypergeometric products)

**SumProd**( $\mathbb{G}$ ) = the smallest set containing  $\mathbb{G} \cup \text{Prod}^*(\mathbb{G})$  with:

1. For all  $f, g \in \text{SumProd}(\mathbb{G})$  we have  $f \oplus g \in \text{SumProd}(\mathbb{G})$ .
2. For all  $f, g \in \text{SumProd}(\mathbb{G})$  we have  $f \odot g \in \text{SumProd}(\mathbb{G})$ .
3. For all  $f \in \text{SumProd}(\mathbb{G})$  and  $k \in \mathbb{Z}_{\geq 1}$  we have  $f^{\Delta k} \in \text{SumProd}(\mathbb{G})$ .
4. For all  $f \in \text{SumProd}(\mathbb{G})$  and  $l \in \mathbb{N}$  we have  $\text{Sum}(l, f) \in \text{SumProd}(\mathbb{G})$ .

Furthermore, the **set of nested sums over hypergeometric products** is given by

$$\Sigma(\mathbb{G}) = \{\text{Sum}(l, f) \mid l \in \mathbb{N} \text{ and } f \in \text{SumProd}(\mathbb{G})\}$$

and the **set of nested sums and hypergeometric products** is given by

$$\Sigma\Pi(\mathbb{G}) = \Sigma(\mathbb{G}) \cup \Pi(\mathbb{G}).$$

$\mathbb{G} \longrightarrow \text{SumProd}(\mathbb{G})$  (nested sums over hypergeometric products)

**SumProd**( $\mathbb{G}$ ) = the smallest set containing  $\mathbb{G} \cup \text{Prod}^*(\mathbb{G})$  with:

1. For all  $f, g \in \text{SumProd}(\mathbb{G})$  we have  $f \oplus g \in \text{SumProd}(\mathbb{G})$ .
2. For all  $f, g \in \text{SumProd}(\mathbb{G})$  we have  $f \odot g \in \text{SumProd}(\mathbb{G})$ .
3. For all  $f \in \text{SumProd}(\mathbb{G})$  and  $k \in \mathbb{Z}_{\geq 1}$  we have  $f^{\Delta k} \in \text{SumProd}(\mathbb{G})$ .
4. For all  $f \in \text{SumProd}(\mathbb{G})$  and  $l \in \mathbb{N}$  we have  $\text{Sum}(l, f) \in \text{SumProd}(\mathbb{G})$ .

### Example

With  $\mathbb{G} = \mathbb{K}(x)$  we get, e.g., the following expressions:

$$E_1 = \text{Sum}(1, \text{Prod}(1, x)) \in \Sigma(\mathbb{G}) \subset \text{SumProd}(\mathbb{G}),$$

$$E_2 = \text{Sum}(1, \frac{1}{x+1} \odot \text{Sum}(1, \frac{1}{x^3}) \odot \text{Sum}(1, \frac{1}{x})) \in \Sigma(\mathbb{G}) \subset \text{SumProd}(\mathbb{G}),$$

$$E_3 = (E_1 \oplus E_2) \odot E_1 \in \text{SumProd}(\mathbb{G}).$$



$$\text{ev} : \mathbb{G} \times \mathbb{N} \rightarrow \mathbb{K} \quad \longrightarrow \quad \text{ev} : \text{SumProd}(\mathbb{G}) \times \mathbb{N} \rightarrow \mathbb{K}$$

$$\text{ev} : \mathbb{G} \times \mathbb{N} \rightarrow \mathbb{K} \quad \longrightarrow \quad \text{ev} : \text{SumProd}(\mathbb{G}) \times \mathbb{N} \rightarrow \mathbb{K}$$

1. For  $f, g \in \text{SumProd}(\mathbb{G})$ ,  $k \in \mathbb{Z} \setminus \{0\}$  ( $k > 0$  if  $f \notin \text{Prod}^*(\mathbb{G})$ ) we set

$$\text{ev}(f^{\Delta k}, n) := \text{ev}(f, n)^k,$$

$$\text{ev}(f \oplus g, n) := \text{ev}(f, n) + \text{ev}(g, n),$$

$$\text{ev}(f \odot g, n) := \text{ev}(f, n) \text{ev}(g, n);$$

$$\text{ev} : \mathbb{G} \times \mathbb{N} \rightarrow \mathbb{K} \quad \longrightarrow \quad \text{ev} : \text{SumProd}(\mathbb{G}) \times \mathbb{N} \rightarrow \mathbb{K}$$

1. For  $f, g \in \text{SumProd}(\mathbb{G})$ ,  $k \in \mathbb{Z} \setminus \{0\}$  ( $k > 0$  if  $f \notin \text{Prod}^*(\mathbb{G})$ ) we set

$$\text{ev}(f^{\Delta} k, n) := \text{ev}(f, n)^k,$$

$$\text{ev}(f \oplus g, n) := \text{ev}(f, n) + \text{ev}(g, n),$$

$$\text{ev}(f \odot g, n) := \text{ev}(f, n) \text{ev}(g, n);$$

2. for  $r \in \mathcal{R}$  and  $\text{Sum}(l, f), \text{Prod}(\lambda, g) \in \text{SumProd}(\mathbb{G})$  we define

$$\text{ev}(\text{RPow}(r), n) := \prod_{i=1}^n r = r^n,$$

$$\text{ev}(\text{Sum}(l, f), n) := \sum_{i=l}^n \text{ev}(f, i),$$

$$\text{ev}(\text{Prod}(\lambda, g), n) := \prod_{i=\lambda}^n \text{ev}(g, i) = \prod_{i=\lambda}^n g(i).$$

$$\text{ev} : \mathbb{G} \times \mathbb{N} \rightarrow \mathbb{K} \quad \longrightarrow \quad \text{ev} : \text{SumProd}(\mathbb{G}) \times \mathbb{N} \rightarrow \mathbb{K}$$

1. For  $f, g \in \text{SumProd}(\mathbb{G})$ ,  $k \in \mathbb{Z} \setminus \{0\}$  ( $k > 0$  if  $f \notin \text{Prod}^*(\mathbb{G})$ ) we set

$$\text{ev}(f^{\Delta} k, n) := \text{ev}(f, n)^k,$$

$$\text{ev}(f \oplus g, n) := \text{ev}(f, n) + \text{ev}(g, n),$$

$$\text{ev}(f \odot g, n) := \text{ev}(f, n) \text{ev}(g, n);$$

2. for  $r \in \mathcal{R}$  and  $\text{Sum}(l, f), \text{Prod}(\lambda, g) \in \text{SumProd}(\mathbb{G})$  we define

$$\text{ev}(\text{RPow}(r), n) := \prod_{i=1}^n r = r^n,$$

$$\text{ev}(\text{Sum}(l, f), n) := \sum_{i=l}^n \text{ev}(f, i),$$

$$\text{ev}(\text{Prod}(\lambda, g), n) := \prod_{i=\lambda}^n \text{ev}(g, i) = \prod_{i=\lambda}^n g(i).$$

Note:  $\Pi(\mathbb{G})$  defines all hypergeometric products (which evaluate to sequences with non-zero entries).

$ev$  applied to  $f \in \text{SumProd}(\mathbb{G})$  represents a sequence.

$f$  can be considered as a simple program and  $ev(f, n)$  with  $n \in \mathbb{N}$  executes it (like an interpreter/compiler) yielding the  $n$ th entry of the represented sequence.

### Definition

For  $F \in \text{SumProd}(\mathbb{G})$  and  $n \in \mathbb{N}$  we write  $F(n) := ev(F, n)$ .

$ev$  applied to  $f \in \text{SumProd}(\mathbb{G})$  represents a sequence.

$f$  can be considered as a simple program and  $ev(f, n)$  with  $n \in \mathbb{N}$  executes it (like an interpreter/compiler) yielding the  $n$ th entry of the represented sequence.

### Definition

For  $F \in \text{SumProd}(\mathbb{G})$  and  $n \in \mathbb{N}$  we write  $F(n) := ev(F, n)$ .

### Example

For  $E_i \in \text{SumProd}(\mathbb{K}(x))$  with  $i = 1, 2, 3$  we get

$$E_1(n) = ev(E_1, n) = ev(\text{Sum}(1, \text{Prod}(1, x)), n) = \sum_{k=1}^n \prod_{i=1}^k i = \sum_{k=1}^n k!,$$

$\text{ev}$  applied to  $f \in \text{SumProd}(\mathbb{G})$  represents a sequence.

$f$  can be considered as a simple program and  $\text{ev}(f, n)$  with  $n \in \mathbb{N}$  executes it (like an interpreter/compiler) yielding the  $n$ th entry of the represented sequence.

### Definition

For  $F \in \text{SumProd}(\mathbb{G})$  and  $n \in \mathbb{N}$  we write  $F(n) := \text{ev}(F, n)$ .

### Example

For  $E_i \in \text{SumProd}(\mathbb{K}(x))$  with  $i = 1, 2, 3$  we get

$$E_1(n) = \text{ev}(E_1, n) = \text{ev}(\text{Sum}(1, \text{Prod}(1, x)), n) = \sum_{k=1}^n \prod_{i=1}^k i = \sum_{k=1}^n k!,$$

$$\begin{aligned} E_2(n) &= \text{ev}(\text{Sum}(1, \frac{1}{x+1} \odot \text{Sum}(1, \frac{1}{x^3}) \odot \text{Sum}(1, \frac{1}{x})), n) \\ &= \sum_{k=1}^n \frac{1}{1+k} \left( \sum_{i=1}^k \frac{1}{i^3} \right) \sum_{i=1}^k \frac{1}{i} \end{aligned}$$

$\text{ev}$  applied to  $f \in \text{SumProd}(\mathbb{G})$  represents a sequence.

$f$  can be considered as a simple program and  $\text{ev}(f, n)$  with  $n \in \mathbb{N}$  executes it (like an interpreter/compiler) yielding the  $n$ th entry of the represented sequence.

### Definition

For  $F \in \text{SumProd}(\mathbb{G})$  and  $n \in \mathbb{N}$  we write  $F(n) := \text{ev}(F, n)$ .

### Example

For  $E_i \in \text{SumProd}(\mathbb{K}(x))$  with  $i = 1, 2, 3$  we get


$$E_1(n) = \text{ev}(E_1, n) = \text{ev}(\text{Sum}(1, \text{Prod}(1, x)), n) = \sum_{k=1}^n \prod_{i=1}^k i = \sum_{k=1}^n k!,$$

$$\begin{aligned} E_2(n) &= \text{ev}(\text{Sum}(1, \frac{1}{x+1} \odot \text{Sum}(1, \frac{1}{x^3}) \odot \text{Sum}(1, \frac{1}{x})), n) \\ &= \sum_{k=1}^n \frac{1}{1+k} \left( \sum_{i=1}^k \frac{1}{i^3} \right) \sum_{i=1}^k \frac{1}{i} \end{aligned}$$

$$E_3(n) = (E_1(n) + E_2(n))E_1(n)$$



## General picture:



## Definition

An expression  $A \in \text{SumProd}(\mathbb{G})$  is in **reduced representation** if

$$A = (f_1 \odot P_1) \oplus (f_2 \odot P_2) \oplus \cdots \oplus (f_r \odot P_r) \quad (4)$$

with  $f_i \in \mathbb{G}^*$  and

$$P_i = (a_{i,1} \overset{\wedge}{z}_{i,1}) \odot (a_{i,2} \overset{\wedge}{z}_{i,2}) \odot \cdots \odot (a_{i,n_i} \overset{\wedge}{z}_{i,n_i})$$

## Definition

An expression  $A \in \text{SumProd}(\mathbb{G})$  is in **reduced representation** if

$$A = (f_1 \odot P_1) \oplus (f_2 \odot P_2) \oplus \cdots \oplus (f_r \odot P_r) \quad (4)$$

with  $f_i \in \mathbb{G}^*$  and

$$P_i = (a_{i,1} \overset{\wedge}{z}_{i,1}) \odot (a_{i,2} \overset{\wedge}{z}_{i,2}) \odot \cdots \odot (a_{i,n_i} \overset{\wedge}{z}_{i,n_i})$$

for  $1 \leq i \leq r$  where

- ▶  $a_{i,j} = \text{Sum}(l_{i,j}, f_{i,j})$  with  $l_{i,j} \in \mathbb{N}$ ,  $f_{i,j} \in \text{SumProd}(\mathbb{G})$  and  $z_{i,j} \in \mathbb{Z}_{\geq 1}$ ,
- ▶  $a_{i,j} = \text{Prod}(l_{i,j}, f_{i,j})$  with  $l_{i,j} \in \mathbb{N}$ ,  $f_{i,j} \in \text{Prod}^*(\mathbb{G})$  and  $z_{i,j} \in \mathbb{Z} \setminus \{0\}$ ,
- ▶  $a_{i,j} = \text{RPow}(f_{i,j})$  with  $f_{i,j} \in \mathcal{R}$  and  $1 \leq z_{i,j} < \text{ord}(r_{i,j})$

such that the following properties hold:

1. for each  $1 \leq i \leq r$  and  $1 \leq j < j' < n_i$  we have  $a_{i,j} \neq a_{i,j'}$ ;
2. for each  $1 \leq i < i' \leq r$  with  $n_i = n_{i'}$  there does not exist a  $\sigma \in S_{n_i}$  with  $P_{i'} = (a_{i,\sigma(1)} \overset{\wedge}{z}_{i,\sigma(1)}) \odot (a_{i,\sigma(2)} \overset{\wedge}{z}_{i,\sigma(2)}) \odot \cdots \odot (a_{i,\sigma(n_i)} \overset{\wedge}{z}_{i,\sigma(n_i)})$ .

## Definition

An expression  $A \in \text{SumProd}(\mathbb{G})$  is in **reduced representation** if

$$A = (f_1 \odot P_1) \oplus (f_2 \odot P_2) \oplus \cdots \oplus (f_r \odot P_r) \quad (4)$$

with  $f_i \in \mathbb{G}^*$

$H \in \text{SumProd}(\mathbb{G})$  is in **sum-product reduced representation** if

- ▶ it is in reduced representation;
- ▶ for each  $\text{Sum}(l, A)$  and  $\text{Prod}(l, A)$  that occur recursively in  $H$  the following holds:
  - ▶  $A$  is in reduced representation as given in (4);
  - ▶  $l \geq \max(L(f_1), \dots, L(f_r))$  (i.e., no poles occur);
  - ▶ the lower bound  $l$  is greater than or equal to the lower bounds of the sums and products inside of  $A$ .

## Definition

An expression  $A \in \text{SumProd}(\mathbb{G})$  is in **reduced representation** if

$$A = (f_1 \odot P_1) \oplus (f_2 \odot P_2) \oplus \cdots \oplus (f_r \odot P_r) \quad (4)$$

with  $f_i \in \mathbb{G}^*$

$H \in \text{SumProd}(\mathbb{G})$  is in **sum-product reduced representation** if

- ▶ it is in reduced representation;
- ▶ for each  $\text{Sum}(l, A)$  and  $\text{Prod}(l, A)$  that occur recursively in  $H$  the following holds:
  - ▶  $A$  is in reduced representation as given in (4);
  - ▶  $l \geq \max(L(f_1), \dots, L(f_r))$  (i.e., no poles occur);
  - ▶ the lower bound  $l$  is greater than or equal to the lower bounds of the sums and products inside of  $A$ .

## Example

$E_3 = (E_1 \oplus E_2) \odot E_1$  is not in reduced representation

## Definition

An expression  $A \in \text{SumProd}(\mathbb{G})$  is in **reduced representation** if

$$A = (f_1 \odot P_1) \oplus (f_2 \odot P_2) \oplus \cdots \oplus (f_r \odot P_r) \quad (4)$$

with  $f_i \in \mathbb{G}^*$

$H \in \text{SumProd}(\mathbb{G})$  is in **sum-product reduced representation** if

- ▶ it is in reduced representation;
- ▶ for each  $\text{Sum}(l, A)$  and  $\text{Prod}(l, A)$  that occur recursively in  $H$  the following holds:
  - ▶  $A$  is in reduced representation as given in (4);
  - ▶  $l \geq \max(L(f_1), \dots, L(f_r))$  (i.e., no poles occur);
  - ▶ the lower bound  $l$  is greater than or equal to the lower bounds of the sums and products inside of  $A$ .

## Example

$E_3 = (E_1 \oplus E_2) \odot E_1$  is not in reduced representation

$\text{Sum}(0, \frac{1}{x})$  is not in sum-product reduced representation

## Definition

An expression  $A \in \text{SumProd}(\mathbb{G})$  is in **reduced representation** if

$$A = (f_1 \odot P_1) \oplus (f_2 \odot P_2) \oplus \cdots \oplus (f_r \odot P_r) \quad (4)$$

with  $f_i \in \mathbb{G}^*$

$H \in \text{SumProd}(\mathbb{G})$  is in **sum-product reduced representation** if

- ▶ it is in reduced representation;
- ▶ for each  $\text{Sum}(l, A)$  and  $\text{Prod}(l, A)$  that occur recursively in  $H$  the following holds:
  - ▶  $A$  is in reduced representation as given in (4);
  - ▶  $l \geq \max(L(f_1), \dots, L(f_r))$  (i.e., no poles occur);
  - ▶ the lower bound  $l$  is greater than or equal to the lower bounds of the sums and products inside of  $A$ .

## Example

$E_3 = (E_1 \oplus E_2) \odot E_1$  is not in reduced representation

$\text{Sum}(0, \frac{1}{x})$  is not in sum-product reduced representation

$\text{Sum}(1, \text{Sum}(2, \frac{1}{x}))$  is not in sum-product reduced representation

## Definition

An expression  $A \in \text{SumProd}(\mathbb{G})$  is in **reduced representation** if

$$A = (f_1 \odot P_1) \oplus (f_2 \odot P_2) \oplus \cdots \oplus (f_r \odot P_r) \quad (4)$$

with  $f_i \in \mathbb{G}^*$

$H \in \text{SumProd}(\mathbb{G})$  is in **sum-product reduced representation** if

- ▶ it is in reduced representation;
- ▶ for each  $\text{Sum}(l, A)$  and  $\text{Prod}(l, A)$  that occur recursively in  $H$  the following holds:
  - ▶  $A$  is in reduced representation as given in (4);
  - ▶  $l \geq \max(L(f_1), \dots, L(f_r))$  (i.e., no poles occur);
  - ▶ the lower bound  $l$  is greater than or equal to the lower bounds of the sums and products inside of  $A$ .

## Lemma

For any  $A \in \text{SumProd}(\mathbb{G})$ , there is a  $B \in \text{SumProd}(\mathbb{G})$  in sum-product reduced representation and  $\lambda \in \mathbb{N}$  such that

$$A(n) = B(n) \quad \forall n \geq \lambda.$$



**Key-Definitions:** Let  $W \subseteq \Sigma\Pi(\mathbb{G})$ .

**SumProd** $(W, \mathbb{G})$  = the set of elements from  $\text{SumProd}(\mathbb{G})$  which are in reduced representation and the arising sums/products are taken from  $W$ .

**Key-Definitions:** Let  $W \subseteq \Sigma\Pi(\mathbb{G})$ .

**SumProd** $(W, \mathbb{G})$  = the set of elements from  $\text{SumProd}(\mathbb{G})$  which are in reduced representation and the arising sums/products are taken from  $W$ .

- ▶  $W$  is called **shift-closed over**  $\mathbb{G}$  if for any  $A \in \text{SumProd}(W, \mathbb{G})$ ,  $s \in \mathbb{Z}$  there are  $B \in \text{SumProd}(W, \mathbb{G})$  and  $\delta \in \mathbb{N}$  such that

$$A(n + s) = B(n) \quad \forall n \geq \delta.$$

**Key-Definitions:** Let  $W \subseteq \Sigma\Pi(\mathbb{G})$ .

**SumProd** $(W, \mathbb{G})$  = the set of elements from  $\text{SumProd}(\mathbb{G})$  which are in reduced representation and the arising sums/products are taken from  $W$ .

- ▶  $W$  is called **shift-closed over**  $\mathbb{G}$  if for any  $A \in \text{SumProd}(W, \mathbb{G})$ ,  $s \in \mathbb{Z}$  there are  $B \in \text{SumProd}(W, \mathbb{G})$  and  $\delta \in \mathbb{N}$  such that

$$A(n + s) = B(n) \quad \forall n \geq \delta.$$

- ▶  $W$  is called **shift-stable over**  $\mathbb{G}$  if for any product or sum in  $W$  the multiplicand or summand is built by sums and products from  $W$ .

**Key-Definitions:** Let  $W \subseteq \Sigma\Pi(\mathbb{G})$ .

**SumProd** $(W, \mathbb{G})$  = the set of elements from  $\text{SumProd}(\mathbb{G})$  which are in reduced representation and the arising sums/products are taken from  $W$ .

- ▶  $W$  is called **shift-closed over**  $\mathbb{G}$  if for any  $A \in \text{SumProd}(W, \mathbb{G})$ ,  $s \in \mathbb{Z}$  there are  $B \in \text{SumProd}(W, \mathbb{G})$  and  $\delta \in \mathbb{N}$  such that

$$A(n + s) = B(n) \quad \forall n \geq \delta.$$

- ▶  $W$  is called **shift-stable over**  $\mathbb{G}$  if for any product or sum in  $W$  the multiplicand or summand is built by sums and products from  $W$ .

### Example

$W = \{\text{Sum}(1, \text{Sum}(1, \frac{1}{x}), \frac{1}{x})\}$  is neither shift-closed nor shift-stable;

**Key-Definitions:** Let  $W \subseteq \Sigma\Pi(\mathbb{G})$ .

**SumProd**( $W, \mathbb{G}$ ) = the set of elements from  $\text{SumProd}(\mathbb{G})$  which are in reduced representation and the arising sums/products are taken from  $W$ .

- ▶  $W$  is called **shift-closed over**  $\mathbb{G}$  if for any  $A \in \text{SumProd}(W, \mathbb{G})$ ,  $s \in \mathbb{Z}$  there are  $B \in \text{SumProd}(W, \mathbb{G})$  and  $\delta \in \mathbb{N}$  such that

$$A(n + s) = B(n) \quad \forall n \geq \delta.$$

- ▶  $W$  is called **shift-stable over**  $\mathbb{G}$  if for any product or sum in  $W$  the multiplicand or summand is built by sums and products from  $W$ .

### Example

$W = \{\text{Sum}(1, \text{Sum}(1, \frac{1}{x}), \frac{1}{x})\}$  is neither shift-closed nor shift-stable;

$W = \{\text{Sum}(1, \frac{1}{x}), \text{Sum}(1, \text{Sum}(1, \frac{1}{x}), \frac{1}{x})\}$  is shift-closed and shift-stable;

**Key-Definitions:** Let  $W \subseteq \Sigma\Pi(\mathbb{G})$ .

**SumProd**( $W, \mathbb{G}$ ) = the set of elements from  $\text{SumProd}(\mathbb{G})$  which are in reduced representation and the arising sums/products are taken from  $W$ .

- ▶  $W$  is called **shift-closed over**  $\mathbb{G}$  if for any  $A \in \text{SumProd}(W, \mathbb{G})$ ,  $s \in \mathbb{Z}$  there are  $B \in \text{SumProd}(W, \mathbb{G})$  and  $\delta \in \mathbb{N}$  such that

$$A(n + s) = B(n) \quad \forall n \geq \delta.$$

- ▶  $W$  is called **shift-stable over**  $\mathbb{G}$  if for any product or sum in  $W$  the multiplicand or summand is built by sums and products from  $W$ .

### Example

$W = \{\text{Sum}(1, \text{Sum}(1, \frac{1}{x}), \frac{1}{x})\}$  is neither shift-closed nor shift-stable;

$W = \{\text{Sum}(1, \frac{1}{x}), \text{Sum}(1, \text{Sum}(1, \frac{1}{x}), \frac{1}{x})\}$  is shift-closed and shift-stable;

$$W \text{ is shift-stable} \quad \begin{array}{c} \Rightarrow \\ \not\Leftarrow \end{array} \quad W \text{ is shift-closed}$$

**Key-Definitions:** Let  $W \subseteq \Sigma\Pi(\mathbb{G})$ .

**SumProd**( $W, \mathbb{G}$ ) = the set of elements from  $\text{SumProd}(\mathbb{G})$  which are in reduced representation and the arising sums/products are taken from  $W$ .

- ▶  $W$  is called **shift-closed over**  $\mathbb{G}$  if for any  $A \in \text{SumProd}(W, \mathbb{G})$ ,  $s \in \mathbb{Z}$  there are  $B \in \text{SumProd}(W, \mathbb{G})$  and  $\delta \in \mathbb{N}$  such that

$$A(n + s) = B(n) \quad \forall n \geq \delta.$$

- ▶  $W$  is called **shift-stable over**  $\mathbb{G}$  if for any product or sum in  $W$  the multiplicand or summand is built by sums and products from  $W$ .
- ▶  $W$  is called **canonical reduced over**  $\mathbb{G}$  if for any  $A, B \in \text{SumProd}(W, \mathbb{G})$  with

$$A(n) = B(n) \quad \forall n \geq \delta$$

for some  $\delta \in \mathbb{N}$  the following holds:  $A$  and  $B$  are the same up to permutations of the operands in  $\oplus$  and  $\odot$ .

## Definition

$W \subseteq \Sigma\Pi(\mathbb{G})$  is called  **$\sigma$ -reduced over  $\mathbb{G}$**  if

1. the elements in  $W$  are in sum-product reduced form,
2.  $W$  is shift-stable (and thus shift-closed) and
3.  $W$  is canonical reduced.

In particular,  $A \in \text{SumProd}(W, \mathbb{G})$  is called  **$\sigma$ -reduced (w.r.t.  $W$ )** if  $W$  is  $\sigma$ -reduced over  $\mathbb{G}$ .



## Definition

$W \subseteq \Sigma\Pi(\mathbb{G})$  is called  **$\sigma$ -reduced over  $\mathbb{G}$**  if

1. the elements in  $W$  are in sum-product reduced form,
2.  $W$  is shift-stable (and thus shift-closed) and
3.  $W$  is canonical reduced.

In particular,  $A \in \text{SumProd}(W, \mathbb{G})$  is called  **$\sigma$ -reduced (w.r.t.  $W$ )** if  $W$  is  $\sigma$ -reduced over  $\mathbb{G}$ .

**Problem SigmaReduce:** Compute a  $\sigma$ -reduced representation

Given:  $A_1, \dots, A_u \in \text{SumProd}(\mathbb{G})$  with  $\mathbb{G} = \mathbb{K}(x)$ .

Find: a  $\sigma$ -reduced set  $W = \{T_1, \dots, T_e\} \subset \Sigma\Pi(\mathbb{G})$ ,  
 $B_1, \dots, B_u \in \text{SumProd}(W, \mathbb{G})$  and  $\delta_1, \dots, \delta_u \in \mathbb{N}$   
 such that for all  $1 \leq i \leq r$  we get

$$A_i(n) = B_i(n) \quad n \geq \delta_i.$$

- **Canonical representation in term algebras**

$$\begin{array}{c} A_1 \\ \downarrow \\ B_1 \end{array}$$

$\sigma$ -reduced

in  $\text{SumProd}(\mathbb{G})$

$$\forall n \geq \delta \quad \text{ev}(A_1, n) = \text{ev}(B_1, n)$$

- **Canonical representation in term algebras**

$$\begin{array}{c} A_1 \\ \downarrow \\ B_1 \end{array}$$

$\sigma$ -reduced

$$\begin{array}{c} A_2 \\ \downarrow \\ B_2 \end{array}$$

in  $\text{SumProd}(\mathbb{G})$

$$\forall n \geq \delta \quad \text{ev}(A_1, n) = \text{ev}(B_1, n)$$

$$\text{ev}(A_2, n) = \text{ev}(B_2, n)$$

- **Canonical representation in term algebras**

$$\begin{array}{c} A_1 \\ \downarrow \\ B_1 \end{array}$$

$\sigma$ -reduced

$$\begin{array}{c} A_2 \\ \downarrow \\ B_2 \end{array}$$

in  $\text{SumProd}(\mathbb{G})$


$$\forall n \geq \delta \quad \text{ev}(A_1, n) = \text{ev}(B_1, n) \quad = \quad \text{ev}(A_2, n) = \text{ev}(B_2, n)$$



canonical simplifier

$$B_1 = B_2$$

## General picture:



## General picture:

Part 1: Symbolic summation (a short introduction)

Part 2: Modeling of sequences with a term algebra (user interface)

Part 3: Modeling of sequences in difference rings (computer algebra)

Part 4: Construction of appropriate difference rings (advanced CA)

Part 5: Applications

Represent  $H = \text{Sum}(1, \frac{1}{x}) \in \text{SumProd}(\mathbb{G})$  with

$$H(n) = H_n = \sum_{k=1}^n \frac{1}{k}.$$

Represent  $H = \text{Sum}(1, \frac{1}{x}) \in \text{SumProd}(\mathbb{G})$  with

$$H(n) = H_n = \sum_{k=1}^n \frac{1}{k}.$$

1. a formal ring  $\mathbb{A} = \underbrace{\underbrace{\mathbb{Q}(x)}_{\text{rat. fu. field}} [s]}_{\text{polynomial ring}}$



Represent  $H = \text{Sum}(1, \frac{1}{x}) \in \text{SumProd}(\mathbb{G})$  with

$$H(n) = H_n = \sum_{k=1}^n \frac{1}{k}.$$

1. a formal ring  $\mathbb{A} = \mathbb{Q}(x)[s]$
2. an evaluation function

$$\begin{aligned} \text{ev}' : \mathbb{Q}(x) \times \mathbb{N} &\rightarrow \mathbb{Q} \\ \left(\frac{p(x)}{q(x)}, n\right) &\mapsto \begin{cases} \frac{p(n)}{q(n)} & \text{if } q(n) \neq 0 \\ 0 & \text{otherwise} \end{cases} \end{aligned}$$

Represent  $H = \text{Sum}(1, \frac{1}{x}) \in \text{SumProd}(\mathbb{G})$  with

$$H(n) = H_n = \sum_{k=1}^n \frac{1}{k}.$$

1. a formal ring  $\mathbb{A} = \mathbb{Q}(x)[s]$
2. an evaluation function

$$\begin{aligned} \text{ev}' : \mathbb{Q}(x) \times \mathbb{N} &\rightarrow \mathbb{Q} \\ \left(\frac{p(x)}{q(x)}, n\right) &\mapsto \begin{cases} \frac{p(n)}{q(n)} & \text{if } q(n) \neq 0 \\ 0 & \text{otherwise} \end{cases} \end{aligned}$$

$$\text{ev} : \mathbb{Q}(x)[s] \times \mathbb{N} \rightarrow \mathbb{Q}$$

$$\text{ev}(s, n) = H_n$$

Represent  $H = \text{Sum}(1, \frac{1}{x}) \in \text{SumProd}(\mathbb{G})$  with

$$H(n) = H_n = \sum_{k=1}^n \frac{1}{k}.$$

1. a formal ring  $\mathbb{A} = \mathbb{Q}(x)[s]$
2. an evaluation function  $\text{ev} : \mathbb{A} \times \mathbb{N} \rightarrow \mathbb{Q}$

$$\begin{aligned} \text{ev}' : \mathbb{Q}(x) \times \mathbb{N} &\rightarrow \mathbb{Q} \\ \left(\frac{p(x)}{q(x)}, n\right) &\mapsto \begin{cases} \frac{p(n)}{q(n)} & \text{if } q(n) \neq 0 \\ 0 & \text{otherwise} \end{cases} \end{aligned}$$

$$\begin{aligned} \text{ev} : \mathbb{Q}(x)[s] \times \mathbb{N} &\rightarrow \mathbb{Q} \\ \left(\sum_{i=0}^d f_i s^i, n\right) &\mapsto \sum_{i=0}^d \text{ev}'(f_i, n) H_n^i \end{aligned} \quad \text{ev}(s, n) = H_n$$

**Definition:**  $(\mathbb{A}, \text{ev})$  is called an eval-ring

Represent  $H = \text{Sum}(1, \frac{1}{x}) \in \text{SumProd}(\mathbb{G})$  with

$$H(n) = H_n = \sum_{k=1}^n \frac{1}{k}.$$

1. a formal ring  $\mathbb{A} = \mathbb{Q}(x)[s]$
2. an evaluation function  $\text{ev} : \mathbb{A} \times \mathbb{N} \rightarrow \mathbb{Q}$

Consider the map

$$\begin{aligned} \tau : \mathbb{A} &\rightarrow \mathbb{Q}^{\mathbb{N}} \\ f &\mapsto \langle \text{ev}(f, n) \rangle_{n \geq 0} \end{aligned}$$

It is **almost** a ring homomorphism :

$$\tau(x)\tau\left(\frac{1}{x}\right) = \langle 0, 1, 2, 3, \dots \rangle \langle 0, 1, \frac{1}{2}, \frac{1}{3}, \dots \rangle$$

Represent  $H = \text{Sum}(1, \frac{1}{x}) \in \text{SumProd}(\mathbb{G})$  with

$$H(n) = H_n = \sum_{k=1}^n \frac{1}{k}.$$

1. a formal ring  $\mathbb{A} = \mathbb{Q}(x)[s]$
2. an evaluation function  $\text{ev} : \mathbb{A} \times \mathbb{N} \rightarrow \mathbb{Q}$

Consider the map

$$\begin{aligned} \tau : \mathbb{A} &\rightarrow \mathbb{Q}^{\mathbb{N}} \\ f &\mapsto \langle \text{ev}(f, n) \rangle_{n \geq 0} \end{aligned}$$

It is **almost** a ring homomorphism :

$$\begin{aligned} \tau(x)\tau\left(\frac{1}{x}\right) &= \langle 0, 1, 2, 3, \dots \rangle \langle 0, 1, \frac{1}{2}, \frac{1}{3}, \dots \rangle \\ &\quad \parallel \\ &= \langle 0, 1, 1, 1, \dots \rangle \end{aligned}$$

Represent  $H = \text{Sum}(1, \frac{1}{x}) \in \text{SumProd}(\mathbb{G})$  with

$$H(n) = H_n = \sum_{k=1}^n \frac{1}{k}.$$

1. a formal ring  $\mathbb{A} = \mathbb{Q}(x)[s]$
2. an evaluation function  $\text{ev} : \mathbb{A} \times \mathbb{N} \rightarrow \mathbb{Q}$

Consider the map

$$\begin{aligned} \tau : \mathbb{A} &\rightarrow \mathbb{Q}^{\mathbb{N}} \\ f &\mapsto \langle \text{ev}(f, n) \rangle_{n \geq 0} \end{aligned}$$

It is **almost** a ring homomorphism :

$$\begin{aligned} \tau(x)\tau\left(\frac{1}{x}\right) &= \langle 0, 1, 2, 3, \dots \rangle \langle 0, 1, \frac{1}{2}, \frac{1}{3}, \dots \rangle \\ &\quad \parallel \\ &= \langle 0, 1, 1, 1, \dots \rangle \\ &\quad \neq \\ \tau\left(x \frac{1}{x}\right) = \tau(1) &= \langle 1, 1, 1, 1, \dots \rangle \end{aligned}$$

Represent  $H = \text{Sum}(1, \frac{1}{x}) \in \text{SumProd}(\mathbb{G})$  with

$$H(n) = H_n = \sum_{k=1}^n \frac{1}{k}.$$

1. a formal ring  $\mathbb{A} = \mathbb{Q}(x)[s]$
2. an evaluation function  $\text{ev} : \mathbb{A} \times \mathbb{N} \rightarrow \mathbb{Q}$

Consider the map

$$\begin{array}{ll} \tau : \mathbb{A} & \rightarrow \mathbb{Q}^{\mathbb{N}} / \sim \\ f & \mapsto \langle \text{ev}(f, n) \rangle_{n \geq 0} \end{array} \quad \begin{array}{l} (a_n) \sim (b_n) \text{ iff } a_n = b_n \\ \text{from a certain point on} \end{array}$$

It is a ring homomorphism :

$$\begin{array}{ll} \tau(x)\tau(\frac{1}{x}) & = \langle 0, 1, 2, 3, \dots \rangle \langle 0, 1, \frac{1}{2}, \frac{1}{3}, \dots \rangle \\ & \quad \parallel \\ & \langle 0, 1, 1, 1, \dots \rangle \\ & \quad \parallel \\ \tau(x \frac{1}{x}) = \tau(1) & = \langle 1, 1, 1, 1, \dots \rangle \end{array}$$

Represent  $H = \text{Sum}(1, \frac{1}{x}) \in \text{SumProd}(\mathbb{G})$  with

$$H(n) = H_n = \sum_{k=1}^n \frac{1}{k}.$$

1. a formal ring  $\mathbb{A} = \mathbb{Q}(x)[s]$
2. an evaluation function  $\text{ev} : \mathbb{A} \times \mathbb{N} \rightarrow \mathbb{Q}$

Consider the map

$$\begin{array}{ll} \tau : \mathbb{A} & \rightarrow \mathbb{Q}^{\mathbb{N}} / \sim \\ f & \mapsto \langle \text{ev}(f, n) \rangle_{n \geq 0} \end{array} \quad \begin{array}{l} (a_n) \sim (b_n) \text{ iff } a_n = b_n \\ \text{from a certain point on} \end{array}$$

It is an **injective** ring homomorphism (**ring embedding**):

$$\begin{array}{ll} \tau(x)\tau(\frac{1}{x}) & = \langle 0, 1, 2, 3, \dots \rangle \langle 0, 1, \frac{1}{2}, \frac{1}{3}, \dots \rangle \\ & \quad \parallel \\ & \quad \langle 0, 1, 1, 1, \dots \rangle \\ & \quad \parallel \\ \tau(x \frac{1}{x}) = \tau(1) & = \langle 1, 1, 1, 1, \dots \rangle \end{array}$$



Represent  $H = \text{Sum}(1, \frac{1}{x}) \in \text{SumProd}(\mathbb{G})$  with

$$H(n) = H_n = \sum_{k=1}^n \frac{1}{k}.$$

1. a formal ring  $\mathbb{A} = \mathbb{Q}(x)[s]$
2. an evaluation function  $\text{ev} : \mathbb{A} \times \mathbb{N} \rightarrow \mathbb{Q}$
3. a ring automorphism

$$\begin{array}{lcl} \sigma' : \mathbb{Q}(x) & \rightarrow & \mathbb{Q}(x) \\ r(x) & \mapsto & r(x+1) \end{array}$$

Represent  $H = \text{Sum}(1, \frac{1}{x}) \in \text{SumProd}(\mathbb{G})$  with

$$H(n) = H_n = \sum_{k=1}^n \frac{1}{k}.$$

1. a formal ring  $\mathbb{A} = \mathbb{Q}(x)[s]$
2. an evaluation function  $\text{ev} : \mathbb{A} \times \mathbb{N} \rightarrow \mathbb{Q}$
3. a ring automorphism

$$\begin{array}{lll} \sigma' : \mathbb{Q}(x) & \rightarrow & \mathbb{Q}(x) \\ & & r(x) \mapsto r(x+1) \end{array}$$

$$\sigma : \mathbb{Q}(x)[s] \rightarrow \mathbb{Q}(x)[s] \qquad s \mapsto s + \frac{1}{x+1}$$

$$H_{n+1} = H_n + \frac{1}{n+1}$$

Represent  $H = \text{Sum}(1, \frac{1}{x}) \in \text{SumProd}(\mathbb{G})$  with

$$H(n) = H_n = \sum_{k=1}^n \frac{1}{k}.$$

1. a formal ring  $\mathbb{A} = \mathbb{Q}(x)[s]$
2. an evaluation function  $\text{ev} : \mathbb{A} \times \mathbb{N} \rightarrow \mathbb{Q}$
3. a ring automorphism

$$\begin{aligned} \sigma' : \mathbb{Q}(x) &\rightarrow \mathbb{Q}(x) \\ r(x) &\mapsto r(x+1) \end{aligned}$$

$$\begin{aligned} \sigma : \mathbb{Q}(x)[s] &\rightarrow \mathbb{Q}(x)[s] & s &\mapsto s + \frac{1}{x+1} \\ \sum_{i=0}^d f_i s^i &\mapsto \sum_{i=0}^d \sigma'(f_i) \left( s + \frac{1}{x+1} \right)^i & H_{n+1} &= H_n + \frac{1}{n+1} \end{aligned}$$

**Definition:**  $(\mathbb{A}, \sigma)$  with a ring  $\mathbb{A}$  and automorphism  $\sigma$  is called a difference ring; the set of constants is

$$\text{const}_\sigma \mathbb{A} = \{c \in \mathbb{A} \mid \sigma(c) = c\}$$

Represent  $H = \text{Sum}(1, \frac{1}{x}) \in \text{SumProd}(\mathbb{G})$  with

$$H(n) = H_n = \sum_{k=1}^n \frac{1}{k}.$$

1. a formal ring  $\mathbb{A} = \mathbb{Q}(x)[s]$
2. an evaluation function  $\text{ev} : \mathbb{A} \times \mathbb{N} \rightarrow \mathbb{Q}$
3. a ring automorphism  $\sigma : \mathbb{A} \rightarrow \mathbb{A}$

ev and  $\sigma$  interact:

$$\text{ev}(\sigma(s), n) = \text{ev}(s + \frac{1}{x+1}, n) = H_n + \frac{1}{n+1} = \text{ev}(s, n+1)$$

Represent  $H = \text{Sum}(1, \frac{1}{x}) \in \text{SumProd}(\mathbb{G})$  with

$$H(n) = H_n = \sum_{k=1}^n \frac{1}{k}.$$

1. a formal ring  $\mathbb{A} = \mathbb{Q}(x)[s]$
2. an evaluation function  $\text{ev} : \mathbb{A} \times \mathbb{N} \rightarrow \mathbb{Q}$
3. a ring automorphism  $\sigma : \mathbb{A} \rightarrow \mathbb{A}$

ev and  $\sigma$  interact:

$$\text{ev}(\sigma(s), n) = \text{ev}(s + \frac{1}{x+1}, n) \Downarrow = H_n + \frac{1}{n+1} = \text{ev}(s, n+1)$$

$$\tau(\sigma(s)) = \langle 1, 1 + \frac{1}{2}, 1 + \frac{1}{2} + \frac{1}{3}, \dots \rangle = S(\langle 0, 1, 1 + \frac{1}{2}, \dots \rangle) = S(\tau(s))$$

shift operator 

Represent  $H = \text{Sum}(1, \frac{1}{x}) \in \text{SumProd}(\mathbb{G})$  with

$$H(n) = H_n = \sum_{k=1}^n \frac{1}{k}.$$

1. a formal ring  $\mathbb{A} = \mathbb{Q}(x)[s]$
2. an evaluation function  $\text{ev} : \mathbb{A} \times \mathbb{N} \rightarrow \mathbb{Q}$
3. a ring automorphism  $\sigma : \mathbb{A} \rightarrow \mathbb{A}$

ev and  $\sigma$  interact:

$$\text{ev}(\sigma(s), n) = \text{ev}(s + \frac{1}{x+1}, n) = H_n + \frac{1}{n+1} = \text{ev}(s, n+1)$$

$\tau(\sigma(s)) = \langle 1, 1 + \frac{1}{2}, 1 + \frac{1}{2} + \frac{1}{3}, \dots \rangle = S(\langle 0, 1, 1 + \frac{1}{2}, \dots \rangle) = S(\tau(s))$   
 $\tau$  is an **injective** difference ring homomorphism:

$$\begin{array}{ccc} \mathbb{K}(x)[s] & \xrightarrow{\sigma} & \mathbb{K}(x)[s] \\ \downarrow \tau & = & \downarrow \tau \\ \mathbb{K}^{\mathbb{N}} / \sim & \xrightarrow{S} & \mathbb{K}^{\mathbb{N}} / \sim \end{array}$$

Represent  $H = \text{Sum}(1, \frac{1}{x}) \in \text{SumProd}(\mathbb{G})$  with

$$H(n) = H_n = \sum_{k=1}^n \frac{1}{k}.$$

1. a formal ring  $\mathbb{A} = \mathbb{Q}(x)[s]$
2. an evaluation function  $\text{ev} : \mathbb{A} \times \mathbb{N} \rightarrow \mathbb{Q}$
3. a ring automorphism  $\sigma : \mathbb{A} \rightarrow \mathbb{A}$

ev and  $\sigma$  interact:

$$\text{ev}(\sigma(s), n) = \text{ev}(s + \frac{1}{x+1}, n) = H_n + \frac{1}{n+1} = \text{ev}(s, n+1)$$

$$\tau(\sigma(s)) = \langle 1, 1 + \frac{1}{2}, 1 + \frac{1}{2} + \frac{1}{3}, \dots \rangle = S(\langle 0, 1, 1 + \frac{1}{2}, \dots \rangle) = S(\tau(s))$$

$\tau$  is an **injective** difference ring homomorphism:

$$\boxed{(\mathbb{K}(x)[s], \sigma)} \stackrel{\tau}{\simeq} \boxed{\underbrace{(\tau(\mathbb{Q}(x))[\langle H_n \rangle_{n \geq 0}], S)}_{\text{rat. seq.}}} \leq (\mathbb{K}^{\mathbb{N}} / \sim, S)$$

Summary: we rephrase  $H \in \text{SumProd}(\mathbb{G})$  as element  $h$  in a formal difference ring. More precisely, we will design

- ▶ a ring  $\mathbb{A}$  with  $\mathbb{A} \supseteq \mathbb{G} \supseteq \mathbb{K}$  in which  $H$  can be represented by  $h \in \mathbb{A}$ ;
- ▶ an evaluation function  $\text{ev} : \mathbb{A} \times \mathbb{N} \rightarrow \mathbb{K}$  such that  $H(n) = \text{ev}(h, n)$  holds for sufficiently large  $n \in \mathbb{N}$ ;
- ▶ a ring automorphism  $\sigma : \mathbb{A} \rightarrow \mathbb{A}$  which models  $H(n+1)$  with  $\sigma(h)$ .



A hypergeometric *APS*-extension of  $(\mathbb{K}(x), \sigma)$  is

- ▶ a ring

$$\mathbb{A} := \mathbb{K}(x)$$

- ▶ with an automorphism where  $\sigma(c) = c$  for all  $c \in \mathbb{K}$  and where

$$\sigma(x) = x + 1$$

A hypergeometric *APS*-extension of  $(\mathbb{K}(x), \sigma)$  is

- ▶ a ring

$$\mathbb{A} := \mathbb{K}(x)$$

- ▶ with an automorphism where  $\sigma(c) = c$  for all  $c \in \mathbb{K}$  and where

$$\sigma(x) = x + 1$$

$$S_k! = (k+1)k!$$

A hypergeometric *APS*-extension of  $(\mathbb{K}(x), \sigma)$  is

- ▶ a ring

$$\mathbb{A} := \mathbb{K}(x)[p_1, p_1^{-1}]$$

- ▶ with an automorphism where  $\sigma(c) = c$  for all  $c \in \mathbb{K}$  and where

$$\sigma(x) = x + 1$$

$$S_k! = (k+1)k! \quad \leftrightarrow \quad \sigma(p_1) = (x+1)p_1$$

A hypergeometric *APS*-extension of  $(\mathbb{K}(x), \sigma)$  is

- ▶ a ring

$$\mathbb{A} := \mathbb{K}(x)[p_1, p_1^{-1}]$$

- ▶ with an automorphism where  $\sigma(c) = c$  for all  $c \in \mathbb{K}$  and where

$$\sigma(x) = x + 1$$

hypergeometric products  $\leftrightarrow \sigma(p_1) = a_1 p_1 \quad a_1 \in \mathbb{K}(x)^*$

A hypergeometric *APS*-extension of  $(\mathbb{K}(x), \sigma)$  is

- ▶ a ring

$$\mathbb{A} := \mathbb{K}(x)[p_1, p_1^{-1}][p_2, p_2^{-1}]$$

- ▶ with an automorphism where  $\sigma(c) = c$  for all  $c \in \mathbb{K}$  and where

$$\sigma(x) = x + 1$$

hypergeometric products	$\leftrightarrow$	$\sigma(p_1) = a_1 p_1$	$a_1 \in \mathbb{K}(x)^*$
		$\sigma(p_2) = a_2 p_2$	$a_2 \in \mathbb{K}(x)^*$

A hypergeometric *APS*-extension of  $(\mathbb{K}(x), \sigma)$  is

- ▶ a ring

$$\mathbb{A} := \mathbb{K}(x)[p_1, p_1^{-1}][p_2, p_2^{-1}] \cdots [p_e, p_e^{-1}]$$

- ▶ with an automorphism where  $\sigma(c) = c$  for all  $c \in \mathbb{K}$  and where

$$\sigma(x) = x + 1$$

hypergeometric products	$\leftrightarrow$	$\sigma(p_1) = a_1 p_1$	$a_1 \in \mathbb{K}(x)^*$
		$\sigma(p_2) = a_2 p_2$	$a_2 \in \mathbb{K}(x)^*$
		$\vdots$	
		$\sigma(p_e) = a_e p_e$	$a_e \in \mathbb{K}(x)^*$

A hypergeometric *APS*-extension of  $(\mathbb{K}(x), \sigma)$  is

► a ring

$$\mathbb{A} := \mathbb{K}(x)[p_1, p_1^{-1}][p_2, p_2^{-1}] \cdots [p_e, p_e^{-1}][z]$$

► with an automorphism where  $\sigma(c) = c$  for all  $c \in \mathbb{K}$  and where

$$\sigma(x) = x + 1$$

hypergeometric products	$\leftrightarrow$	$\sigma(p_1) = a_1 p_1$	$a_1 \in \mathbb{K}(x)^*$
		$\sigma(p_2) = a_2 p_2$	$a_2 \in \mathbb{K}(x)^*$
		$\vdots$	
		$\sigma(p_e) = a_e p_e$	$a_e \in \mathbb{K}(x)^*$
$(-1)^k$	$\leftrightarrow$	$\sigma(z) = -z$	$z^2 = 1$

A hypergeometric *APS*-extension of  $(\mathbb{K}(x), \sigma)$  is

► a ring

$$\mathbb{A} := \mathbb{K}(x)[p_1, p_1^{-1}][p_2, p_2^{-1}] \cdots [p_e, p_e^{-1}][z]$$

► with an automorphism where  $\sigma(c) = c$  for all  $c \in \mathbb{K}$  and where

$$\sigma(x) = x + 1$$

$$\begin{array}{l} \text{hypergeometric} \\ \text{products} \end{array} \quad \leftrightarrow \quad \begin{array}{l} \sigma(p_1) = a_1 p_1 \\ \sigma(p_2) = a_2 p_2 \\ \vdots \\ \sigma(p_e) = a_e p_e \end{array} \quad \begin{array}{l} a_1 \in \mathbb{K}(x)^* \\ a_2 \in \mathbb{K}(x)^* \\ \vdots \\ a_e \in \mathbb{K}(x)^* \end{array}$$

$\gamma$  is a primitive  $\lambda$ th  
root of unity

$\gamma^k$

$$\leftrightarrow \quad \sigma(\mathbf{z}) = \gamma \mathbf{z} \quad \mathbf{z}^\lambda = \mathbf{1}$$



A hypergeometric *APS*-extension of  $(\mathbb{K}(x), \sigma)$  is

► a ring

$$\mathbb{A} := \mathbb{K}(x)[p_1, p_1^{-1}][p_2, p_2^{-1}] \cdots [p_e, p_e^{-1}][z][s_1]$$

► with an automorphism where  $\sigma(c) = c$  for all  $c \in \mathbb{K}$  and where

$$\sigma(x) = x + 1$$

$$\begin{array}{l} \text{hypergeometric} \\ \text{products} \end{array} \quad \Leftrightarrow \quad \begin{array}{l} \sigma(p_1) = a_1 p_1 \\ \sigma(p_2) = a_2 p_2 \end{array} \quad \begin{array}{l} a_1 \in \mathbb{K}(x)^* \\ a_2 \in \mathbb{K}(x)^* \end{array}$$

$$\vdots$$

$$\sigma(p_e) = a_e p_e \quad a_e \in \mathbb{K}(x)^*$$

$$\begin{array}{l} \gamma \text{ is a primitive } \lambda\text{th} \\ \text{root of unity} \end{array} \quad \gamma^{\mathbf{k}} \quad \Leftrightarrow \quad \begin{array}{l} \sigma(\mathbf{z}) = \gamma \mathbf{z} \\ \mathbf{z}^\lambda = \mathbf{1} \end{array}$$

$$H_{k+1} = H_k + \frac{1}{k+1} \quad \Leftrightarrow \quad \sigma(s_1) = s_1 + \frac{1}{x+1}$$

A hypergeometric *APS*-extension of  $(\mathbb{K}(x), \sigma)$  is

► a ring

$$\mathbb{A} := \mathbb{K}(x)[p_1, p_1^{-1}][p_2, p_2^{-1}] \cdots [p_e, p_e^{-1}][z][s_1]$$

► with an automorphism where  $\sigma(c) = c$  for all  $c \in \mathbb{K}$  and where

$$\sigma(x) = x + 1$$

$$\begin{array}{l} \text{hypergeometric} \\ \text{products} \end{array} \quad \Leftrightarrow \quad \begin{array}{ll} \sigma(p_1) = a_1 p_1 & a_1 \in \mathbb{K}(x)^* \\ \sigma(p_2) = a_2 p_2 & a_2 \in \mathbb{K}(x)^* \\ \vdots & \\ \sigma(p_e) = a_e p_e & a_e \in \mathbb{K}(x)^* \end{array}$$

$$\begin{array}{l} \gamma \text{ is a primitive } \lambda\text{th} \\ \text{root of unity} \end{array} \quad \gamma^k \quad \Leftrightarrow \quad \begin{array}{ll} \sigma(\mathbf{z}) = \gamma \mathbf{z} & \mathbf{z}^\lambda = \mathbf{1} \end{array}$$

$$\text{(nested) sum} \quad \Leftrightarrow \quad \sigma(s_1) = s_1 + f_1 \quad f_1 \in \mathbb{K}(x)[p_1, p_1^{-1}] \cdots [p_e, p_e^{-1}][z]$$

A hypergeometric  $APS$ -extension of  $(\mathbb{K}(x), \sigma)$  is

► a ring

$$\mathbb{A} := \mathbb{K}(x)[p_1, p_1^{-1}][p_2, p_2^{-1}] \cdots [p_e, p_e^{-1}][z][s_1][s_2]$$

► with an automorphism where  $\sigma(c) = c$  for all  $c \in \mathbb{K}$  and where

$$\sigma(x) = x + 1$$

$$\begin{array}{l} \text{hypergeometric} \\ \text{products} \end{array} \quad \Leftrightarrow \quad \begin{array}{ll} \sigma(p_1) = a_1 p_1 & a_1 \in \mathbb{K}(x)^* \\ \sigma(p_2) = a_2 p_2 & a_2 \in \mathbb{K}(x)^* \end{array}$$

$$\vdots$$

$$\sigma(p_e) = a_e p_e \quad a_e \in \mathbb{K}(x)^*$$

$$\begin{array}{l} \gamma \text{ is a primitive } \lambda\text{th} \\ \text{root of unity} \end{array} \quad \gamma^k \quad \Leftrightarrow \quad \begin{array}{ll} \sigma(\mathbf{z}) = \gamma \mathbf{z} & \mathbf{z}^\lambda = \mathbf{1} \end{array}$$

$$\begin{array}{l} \text{(nested) sum} \end{array} \quad \Leftrightarrow \quad \begin{array}{ll} \sigma(s_1) = s_1 + f_1 & f_1 \in \mathbb{K}(x)[p_1, p_1^{-1}] \cdots [p_e, p_e^{-1}][z] \\ \sigma(s_2) = s_2 + f_2 & f_2 \in \mathbb{K}(x)[p_1, p_1^{-1}] \cdots [p_e, p_e^{-1}][z][s_1] \end{array}$$

A hypergeometric *APS*-extension of  $(\mathbb{K}(x), \sigma)$  is

- ▶ a ring

$$\mathbb{A} := \mathbb{K}(x)[p_1, p_1^{-1}][p_2, p_2^{-1}] \cdots [p_e, p_e^{-1}][z][s_1][s_2][s_3] \cdots$$

- ▶ with an automorphism where  $\sigma(c) = c$  for all  $c \in \mathbb{K}$  and where

$$\sigma(x) = x + 1$$

$$\begin{array}{l} \text{hypergeometric} \\ \text{products} \end{array} \quad \Leftrightarrow \quad \begin{array}{ll} \sigma(p_1) = a_1 p_1 & a_1 \in \mathbb{K}(x)^* \\ \sigma(p_2) = a_2 p_2 & a_2 \in \mathbb{K}(x)^* \end{array}$$

$$\vdots$$

$$\sigma(p_e) = a_e p_e \quad a_e \in \mathbb{K}(x)^*$$

$$\begin{array}{l} \gamma \text{ is a primitive } \lambda\text{th} \\ \text{root of unity} \end{array} \quad \gamma^k \quad \Leftrightarrow \quad \begin{array}{ll} \sigma(\mathbf{z}) = \gamma \mathbf{z} & \mathbf{z}^\lambda = \mathbf{1} \end{array}$$

$$\begin{array}{l} \text{(nested) sum} \end{array} \quad \Leftrightarrow \quad \begin{array}{ll} \sigma(s_1) = s_1 + f_1 & f_1 \in \mathbb{K}(x)[p_1, p_1^{-1}] \cdots [p_e, p_e^{-1}][z] \\ \sigma(s_2) = s_2 + f_2 & f_2 \in \mathbb{K}(x)[p_1, p_1^{-1}] \cdots [p_e, p_e^{-1}][z][s_1] \\ \sigma(s_3) = s_3 + f_3 & f_3 \in \mathbb{K}(x)[p_1, p_1^{-1}] \cdots [p_e, p_e^{-1}][z][s_1][s_2] \\ \vdots & \end{array}$$

## Definition (Evaluation function)

Take  $(\mathbb{A}, \sigma)$  with a subfield  $\mathbb{K}$  of  $\mathbb{A}$  with  $\sigma|_{\mathbb{K}} = \text{id}$ .

1.  $\text{ev} : \mathbb{A} \times \mathbb{N} \rightarrow \mathbb{K}$  is called **evaluation function** for  $(\mathbb{A}, \sigma)$  if for all  $f, g \in \mathbb{A}$ ,  $c \in \mathbb{K}$  and  $l \in \mathbb{Z}$  there exists a  $\lambda \in \mathbb{N}$  with

$$\forall n \geq \lambda : \text{ev}(c, n) = c, \quad (5)$$

$$\forall n \geq \lambda : \text{ev}(f + g, n) = \text{ev}(f, n) + \text{ev}(g, n), \quad (6)$$

$$\forall n \geq \lambda : \text{ev}(f g, n) = \text{ev}(f, n) \text{ev}(g, n), \quad (7)$$

$$\forall n \geq \lambda : \text{ev}(\sigma^l(f), n) = \text{ev}(f, n + l). \quad (8)$$

## Definition (Evaluation function)

Take  $(\mathbb{A}, \sigma)$  with a subfield  $\mathbb{K}$  of  $\mathbb{A}$  with  $\sigma|_{\mathbb{K}} = \text{id}$ .

1.  $\text{ev} : \mathbb{A} \times \mathbb{N} \rightarrow \mathbb{K}$  is called **evaluation function** for  $(\mathbb{A}, \sigma)$  if for all  $f, g \in \mathbb{A}$ ,  $c \in \mathbb{K}$  and  $l \in \mathbb{Z}$  there exists a  $\lambda \in \mathbb{N}$  with

$$\forall n \geq \lambda : \text{ev}(c, n) = c, \quad (5)$$

$$\forall n \geq \lambda : \text{ev}(f + g, n) = \text{ev}(f, n) + \text{ev}(g, n), \quad (6)$$

$$\forall n \geq \lambda : \text{ev}(f g, n) = \text{ev}(f, n) \text{ev}(g, n), \quad (7)$$

$$\forall n \geq \lambda : \text{ev}(\sigma^l(f), n) = \text{ev}(f, n + l). \quad (8)$$

2.  $L : \mathbb{A} \rightarrow \mathbb{N}$  is called ***o*-function** if for any  $f, g \in \mathbb{A}$  with  $\lambda = \max(L(f), L(g))$  the properties (6) and (7) hold and for any  $f \in \mathbb{A}$  and  $l \in \mathbb{Z}$  with  $\lambda = L(f) + \max(0, -l)$  property (8) holds.

Connection between  $\text{SumProd}(\mathbb{G})$  and hypergeometric  $APS$ -extension

- **Observation 1:** Given  $\{T_1, \dots, T_e\} \subseteq \Sigma\Pi(\mathbb{G})$ , one can construct a hypergeometric  $APS$ -extension  $(\mathbb{E}, \sigma)$  of  $(\mathbb{G}, \sigma)$  with  $\text{ev}$  and  $L$  such that there are  $a_1, \dots, a_e \in \mathbb{E}$  and  $\delta_1, \dots, \delta_e$  with  $\text{ev}(a_i, n) = T_i(n)$ .

## Connection between $\text{SumProd}(\mathbb{G})$ and hypergeometric $APS$ -extension

• **Observation 1:** Given  $\{T_1, \dots, T_e\} \subseteq \Sigma\Pi(\mathbb{G})$ , one can construct a hypergeometric  $APS$ -extension  $(\mathbb{E}, \sigma)$  of  $(\mathbb{G}, \sigma)$  with  $\text{ev}$  and  $L$  such that there are  $a_1, \dots, a_e \in \mathbb{E}$  and  $\delta_1, \dots, \delta_e$  with  $\text{ev}(a_i, n) = T_i(n)$ .

• **Observation 2:**

$(\mathbb{E}, \sigma)$  with  $\mathbb{E} = \mathbb{G}\langle t_1 \rangle \dots \langle t_e \rangle$  a hypergeometric  $APS$ -extension of  $(\mathbb{G}, \sigma)$   
 $\text{ev} : \mathbb{E} \times \mathbb{N} \rightarrow \mathbb{K}$ ,  $L : \mathbb{E} \rightarrow \mathbb{N}$



$\forall n \geq L(t_i) :$   
 $\text{ev}(t_i, n) = T_i(n) \in \Sigma\Pi(\mathbb{G})$

$W = \{T_1, \dots, T_e\} \subseteq \Sigma\Pi(\mathbb{G})$  is sum-product reduced and  
**shift stable:** sums/products in  $T_i$  are from  $\{T_1, \dots, T_{i-1}\}$ .

In particular, if  $f \in \mathbb{E} \setminus \{0\}$ , then we can take the "unique"  
 $0 \neq F \in \text{SumProd}(\{T_1, \dots, T_e\}, \mathbb{G})$  with  $F(n) = \text{ev}(f, n)$  for all  $n \geq L(f)$ .



## Connection between $\text{SumProd}(\mathbb{G})$ and hypergeometric $APS$ -extension

• **Observation 1:** Given  $\{T_1, \dots, T_e\} \subseteq \Sigma\Pi(\mathbb{G})$ , one can construct a hypergeometric  $APS$ -extension  $(\mathbb{E}, \sigma)$  of  $(\mathbb{G}, \sigma)$  with  $\text{ev}$  and  $L$  such that there are  $a_1, \dots, a_e \in \mathbb{E}$  and  $\delta_1, \dots, \delta_e$  with  $\text{ev}(a_i, n) = T_i(n)$ .

• **Observation 2:**

$(\mathbb{E}, \sigma)$  with  $\mathbb{E} = \mathbb{G}\langle t_1 \rangle \dots \langle t_e \rangle$  a hypergeometric  $APS$ -extension of  $(\mathbb{G}, \sigma)$   
 $\text{ev} : \mathbb{E} \times \mathbb{N} \rightarrow \mathbb{K}$ ,  $L : \mathbb{E} \rightarrow \mathbb{N}$



$$\forall n \geq L(t_i) : \\ \text{ev}(t_i, n) = T_i(n) \in \Sigma\Pi(\mathbb{G})$$

$W = \{T_1, \dots, T_e\} \subseteq \Sigma\Pi(\mathbb{G})$  is sum-product reduced and  
**shift stable:** sums/products in  $T_i$  are from  $\{T_1, \dots, T_{i-1}\}$ .

In particular, if  $f \in \mathbb{E} \setminus \{0\}$ , then we can take the "unique"  
 $0 \neq F \in \text{SumProd}(\{T_1, \dots, T_e\}, \mathbb{G})$  with  $F(n) = \text{ev}(f, n)$  for all  $n \geq L(f)$ .

### Definition

For  $f \in \mathbb{E}$  we also write  $\text{expr}(f) = F$  for this particular  $F$ .

## Connection between $\text{SumProd}(\mathbb{G})$ and hypergeometric $APS$ -extension

• **Observation 1:** Given  $\{T_1, \dots, T_e\} \subseteq \Sigma\Pi(\mathbb{G})$ , one can construct a hypergeometric  $APS$ -extension  $(\mathbb{E}, \sigma)$  of  $(\mathbb{G}, \sigma)$  with  $\text{ev}$  and  $L$  such that there are  $a_1, \dots, a_e \in \mathbb{E}$  and  $\delta_1, \dots, \delta_e$  with  $\text{ev}(a_i, n) = T_i(n)$ .

• **Observation 2:**

$(\mathbb{E}, \sigma)$  with  $\mathbb{E} = \mathbb{G}\langle t_1 \rangle \dots \langle t_e \rangle$  a hypergeometric  $APS$ -extension of  $(\mathbb{G}, \sigma)$   
 $\text{ev} : \mathbb{E} \times \mathbb{N} \rightarrow \mathbb{K}$ ,  $L : \mathbb{E} \rightarrow \mathbb{N}$



$$\forall n \geq L(t_i) : \\ \text{ev}(t_i, n) = T_i(n) \in \Sigma\Pi(\mathbb{G})$$

$W = \{T_1, \dots, T_e\} \subseteq \Sigma\Pi(\mathbb{G})$  is sum-product reduced and  
**shift stable:** sums/products in  $T_i$  are from  $\{T_1, \dots, T_{i-1}\}$ .

### Example

For  $f = x + \frac{x+1}{x}s^4 \in \mathbb{Q}(x)[s]$  we obtain

$$\text{expr}(f) = F = x \oplus \left(\frac{x+1}{x}\right) \odot (\text{Sum}(1, \frac{1}{x})^{\Delta} 4) \in \text{Sum}(\mathbb{Q}(x))$$

with  $F(n) = \text{ev}(f, n)$  for all  $n \geq 1$ .

## Connection between $\text{SumProd}(\mathbb{G})$ and hypergeometric $APS$ -extension

• **Observation 1:** Given  $\{T_1, \dots, T_e\} \subseteq \Sigma\Pi(\mathbb{G})$ , one can construct a hypergeometric  $APS$ -extension  $(\mathbb{E}, \sigma)$  of  $(\mathbb{G}, \sigma)$  with  $\text{ev}$  and  $L$  such that there are  $a_1, \dots, a_e \in \mathbb{E}$  and  $\delta_1, \dots, \delta_e$  with  $\text{ev}(a_i, n) = T_i(n)$ .

• **Observation 2:**

$(\mathbb{E}, \sigma)$  with  $\mathbb{E} = \mathbb{G}\langle t_1 \rangle \dots \langle t_e \rangle$  a hypergeometric  $APS$ -extension of  $(\mathbb{G}, \sigma)$   
 $\text{ev} : \mathbb{E} \times \mathbb{N} \rightarrow \mathbb{K}$ ,  $L : \mathbb{E} \rightarrow \mathbb{N}$

$$\begin{array}{ccc} \downarrow & \forall n \geq L(t_i) : & \uparrow \\ & \text{ev}(t_i, n) = T_i(n) \in \Sigma\Pi(\mathbb{G}) & \end{array}$$

$W = \{T_1, \dots, T_e\} \subseteq \Sigma\Pi(\mathbb{G})$  is sum-product reduced and **shift stable**: sums/products in  $T_i$  are from  $\{T_1, \dots, T_{i-1}\}$ .

## Difference ring theory in action

Let  $(\mathbb{E}, \sigma)$  be a hypergeometric *APS*-extension of  $(\mathbb{G}, \sigma)$  with  $\text{ev} : \mathbb{E} \times \mathbb{N} \rightarrow \mathbb{K}$  and let  $\tau : \mathbb{E} \rightarrow \mathbb{K}^{\mathbb{N}} / \sim$  be the  $\mathbb{K}$ -homomorphism given by

$$\tau(f) = (\text{ev}(f, n))_{n \geq 0}.$$

## Difference ring theory in action

Let  $(\mathbb{E}, \sigma)$  be a hypergeometric APS-extension of  $(\mathbb{G}, \sigma)$  with  $\text{ev} : \mathbb{E} \times \mathbb{N} \rightarrow \mathbb{K}$  and let  $\tau : \mathbb{E} \rightarrow \mathbb{K}^{\mathbb{N}} / \sim$  be the  $\mathbb{K}$ -homomorphism given by

$$\tau(f) = (\text{ev}(f, n))_{n \geq 0}.$$

### Lemma

Let  $W = \{T_1, \dots, T_e\} \in \Sigma\Pi(\mathbb{G})$  with  $T_i = \text{expr}(t_i)$ . Then:

*W is canonical reduced*  $\Leftrightarrow$   *$\tau$  is injective.*

## Difference ring theory in action

Let  $(\mathbb{E}, \sigma)$  be a hypergeometric APS-extension of  $(\mathbb{G}, \sigma)$  with  $\text{ev} : \mathbb{E} \times \mathbb{N} \rightarrow \mathbb{K}$  and let  $\tau : \mathbb{E} \rightarrow \mathbb{K}^{\mathbb{N}} / \sim$  be the  $\mathbb{K}$ -homomorphism given by

$$\tau(f) = (\text{ev}(f, n))_{n \geq 0}.$$

### Lemma

Let  $W = \{T_1, \dots, T_e\} \in \Sigma\Pi(\mathbb{G})$  with  $T_i = \text{expr}(t_i)$ . Then:

$$W \text{ is canonical reduced} \quad \Leftrightarrow \quad \tau \text{ is injective.}$$

Using difference ring theory we get the following crucial property:

### Theorem

$$\tau \text{ is injective} \quad \Leftrightarrow \quad \text{const}_{\sigma}\mathbb{E} = \mathbb{K}.$$

## Example

For our difference field  $\mathbb{G} = \mathbb{K}(x)$  with  $\sigma(x) = x + 1$  and  $\text{const}_\sigma \mathbb{K} = \mathbb{K}$  we have  $\text{const}_\sigma \mathbb{K}(x) = \mathbb{K}$ .

## Example

For our difference field  $\mathbb{G} = \mathbb{K}(x)$  with  $\sigma(x) = x + 1$  and  $\text{const}_\sigma \mathbb{K} = \mathbb{K}$  we have  $\text{const}_\sigma \mathbb{K}(x) = \mathbb{K}$ .

## Definition

A hypergeometric *APS*-extension  $(\mathbb{E}, \sigma)$  of  $(\mathbb{G}, \sigma)$  is called **hypergeometric *RΠΣ*-extension** if

$$\text{const}_\sigma \mathbb{E} = \mathbb{K}.$$



## Example

For our difference field  $\mathbb{G} = \mathbb{K}(x)$  with  $\sigma(x) = x + 1$  and  $\text{const}_\sigma \mathbb{K} = \mathbb{K}$  we have  $\text{const}_\sigma \mathbb{K}(x) = \mathbb{K}$ .

## Definition

A hypergeometric *APS*-extension  $(\mathbb{E}, \sigma)$  of  $(\mathbb{G}, \sigma)$  is called **hypergeometric  $R\Pi\Sigma$ -extension** if

$$\text{const}_\sigma \mathbb{E} = \mathbb{K}.$$

## Theorem

Let  $W = \{T_1, \dots, T_e\} \subset \Sigma\Pi(\mathbb{G})$  be in sum-product reduced representation and shift-stable, i.e., for each  $1 \leq i \leq e$  the arising sums and products in  $T_i$  are contained in  $\{T_1, \dots, T_{i-1}\}$ . Then the following is equivalent:

1. There is a hypergeometric  $R\Pi\Sigma$ -extension  $(\mathbb{E}, \sigma)$  of  $(\mathbb{G}, \sigma)$  with  $\mathbb{E} = \mathbb{G}\langle t_1 \rangle \dots \langle t_e \rangle$  equipped with an evaluation function  $\text{ev}$  with  $T_i = \text{expr}(t_i) \in \Sigma\Pi(\mathbb{G})$  for  $1 \leq i \leq e$ .
2.  $W$  is  $\sigma$ -reduced over  $\mathbb{G}$ .

This yields a strategy (actually the only strategy for shift-stable sets).

## A Strategy to solve Problem SigmaReduce

Given:  $A_1, \dots, A_u \in \text{SumProd}(\mathbb{G})$  with  $\mathbb{G} = \mathbb{K}(x)$ .


Find: a  $\sigma$ -reduced set  $W = \{T_1, \dots, T_e\} \subset \Sigma\Pi(\mathbb{G})$  with  $B_1, \dots, B_u \in \text{SumProd}(W, \mathbb{G})$  and  $\delta_1, \dots, \delta_u \in \mathbb{N}$  such that  $A_i(n) = B_i(n)$  holds for all  $n \geq \delta_i$  and  $1 \leq i \leq r$ .

1. Construct  $R\Pi\Sigma$ -extension  $(\mathbb{E}, \sigma)$  of  $(\mathbb{G}, \sigma)$  with  $\mathbb{E} = \mathbb{G}\langle t_1 \rangle \dots \langle t_e \rangle$  equipped with  $\text{ev} : \mathbb{E} \times \mathbb{N} \rightarrow \mathbb{K}$  such that we get  $a_1, \dots, a_u \in \mathbb{E}$  and  $\delta_1, \dots, \delta_u \in \mathbb{N}$  with

$$A_i(n) = \text{ev}(a_i, n) \quad \forall n \geq \delta_i. \quad (12)$$

2. Set  $W = \{T_1, \dots, T_e\}$  with  $T_i := \text{expr}(t_i) \in \Sigma\Pi(\mathbb{G})$  for  $1 \leq i \leq e$ .
3. Set  $B_i := \text{expr}(a_i) \in \text{SumProd}(W, \mathbb{G})$  for  $1 \leq i \leq u$ .
4. Return  $W, (B_1, \dots, B_u)$  and  $(\delta_1, \dots, \delta_u)$ .

## General picture:



## General picture:

Part 1: Symbolic summation (a short introduction)

Part 2: Modeling of sequences with a term algebra (user interface)

Part 3: Modeling of sequences in difference rings (computer algebra)

Part 4: Construction of appropriate difference rings (advanced CA)

Part 5: Applications

A hypergeometric *APS*-extension of  $(\mathbb{K}(x), \sigma)$  is

- ▶ a ring

$$\mathbb{A} := \mathbb{K}(x)[p_1, p_1^{-1}][p_2, p_2^{-1}] \cdots [p_e, p_e^{-1}][z][s_1][s_2][s_3] \cdots$$

- ▶ with an automorphism where  $\sigma(c) = c$  for all  $c \in \mathbb{K}$  and where

$$\sigma(x) = x + 1$$

$$\begin{array}{l} \text{hypergeometric} \\ \text{products} \end{array} \quad \Leftrightarrow \quad \begin{array}{ll} \sigma(p_1) = a_1 p_1 & a_1 \in \mathbb{K}(x)^* \\ \sigma(p_2) = a_2 p_2 & a_2 \in \mathbb{K}(x)^* \end{array}$$

$$\vdots$$

$$\sigma(p_e) = a_e p_e \quad a_e \in \mathbb{K}(x)^*$$

$$\begin{array}{l} \gamma \text{ is a primitive } \lambda\text{th} \\ \text{root of unity} \end{array} \quad \gamma^k \quad \Leftrightarrow \quad \begin{array}{ll} \sigma(\mathbf{z}) = \gamma \mathbf{z} & \mathbf{z}^\lambda = \mathbf{1} \end{array}$$

$$\begin{array}{l} \text{(nested) sum} \end{array} \quad \Leftrightarrow \quad \begin{array}{ll} \sigma(s_1) = s_1 + f_1 & f_1 \in \mathbb{K}(x)[p_1, p_1^{-1}] \cdots [p_e, p_e^{-1}][z] \\ \sigma(s_2) = s_2 + f_2 & f_2 \in \mathbb{K}(x)[p_1, p_1^{-1}] \cdots [p_e, p_e^{-1}][z][s_1] \\ \sigma(s_3) = s_3 + f_3 & f_3 \in \mathbb{K}(x)[p_1, p_1^{-1}] \cdots [p_e, p_e^{-1}][z][s_1][s_2] \\ \vdots & \end{array}$$

A hypergeometric *RHS-extension* of  $(\mathbb{K}(x), \sigma)$  is

- ▶ a ring

$$\mathbb{A} := \mathbb{K}(x)[p_1, p_1^{-1}][p_2, p_2^{-1}] \cdots [p_e, p_e^{-1}][z][s_1][s_2][s_3] \cdots$$

- ▶ with an automorphism where  $\sigma(c) = c$  for all  $c \in \mathbb{K}$  and where

$$\sigma(x) = x + 1$$

$$\begin{array}{l} \text{hypergeometric} \\ \text{products} \end{array} \quad \Leftrightarrow \quad \begin{array}{ll} \sigma(p_1) = a_1 p_1 & a_1 \in \mathbb{K}(x)^* \\ \sigma(p_2) = a_2 p_2 & a_2 \in \mathbb{K}(x)^* \end{array}$$

$$\vdots$$

$$\sigma(p_e) = a_e p_e \quad a_e \in \mathbb{K}(x)^*$$

$$\begin{array}{l} \gamma \text{ is a primitive } \lambda\text{th} \\ \text{root of unity} \end{array} \quad \gamma^k \quad \Leftrightarrow \quad \begin{array}{ll} \sigma(\mathbf{z}) = \gamma \mathbf{z} & \mathbf{z}^\lambda = \mathbf{1} \end{array}$$

$$\begin{array}{l} \text{(nested) sum} \end{array} \quad \Leftrightarrow \quad \begin{array}{ll} \sigma(s_1) = s_1 + f_1 & f_1 \in \mathbb{K}(x)[p_1, p_1^{-1}] \cdots [p_e, p_e^{-1}][z] \\ \sigma(s_2) = s_2 + f_2 & f_2 \in \mathbb{K}(x)[p_1, p_1^{-1}] \cdots [p_e, p_e^{-1}][z][s_1] \\ \sigma(s_3) = s_3 + f_3 & f_3 \in \mathbb{K}(x)[p_1, p_1^{-1}] \cdots [p_e, p_e^{-1}][z][s_1][s_2] \\ \vdots & \end{array}$$

such that  $\text{const}_\sigma \mathbb{E} = \mathbb{K}$

## Represent sums (extension of Karr's result, 1981)

- ▶ Let  $(\mathbb{A}, \sigma)$  be a difference ring with constant set

$$\text{const}_\sigma \mathbb{A} := \{k \in \mathbb{A} \mid \sigma(k) = k\}.$$

**Note 1:**  $\text{const}_\sigma \mathbb{A}$  is a ring that contains  $\mathbb{Q}$

**Note 2:** We always take care that  $\text{const}_\sigma \mathbb{A}$  is a field

## Represent sums (extension of Karr's result, 1981)

- ▶ Let  $(\mathbb{A}, \sigma)$  be a difference ring with constant field

$$\text{const}_\sigma \mathbb{A} := \{k \in \mathbb{A} \mid \sigma(k) = k\}.$$

- ▶ Adjoin a new variable  $t$  to  $\mathbb{A}$  (i.e.,  $\mathbb{A}[t]$  is a polynomial ring).



## Represent sums (extension of Karr's result, 1981)

- ▶ Let  $(\mathbb{A}, \sigma)$  be a difference ring with constant field

$$\text{const}_\sigma \mathbb{A} := \{k \in \mathbb{A} \mid \sigma(k) = k\}.$$

- ▶ Adjoin a new variable  $t$  to  $\mathbb{A}$  (i.e.,  $\mathbb{A}[t]$  is a polynomial ring).
- ▶ Extend the shift operator s.t.

$$\sigma(t) = t + f \quad \text{for some } f \in \mathbb{A}.$$

## Represent sums (extension of Karr's result, 1981)

- ▶ Let  $(\mathbb{A}, \sigma)$  be a difference ring with constant field

$$\text{const}_\sigma \mathbb{A} := \{k \in \mathbb{A} \mid \sigma(k) = k\}.$$

- ▶ Adjoin a new variable  $t$  to  $\mathbb{A}$  (i.e.,  $\mathbb{A}[t]$  is a polynomial ring).
- ▶ Extend the shift operator s.t.

$$\sigma(t) = t + f \quad \text{for some } f \in \mathbb{A}.$$

Then  $\text{const}_\sigma \mathbb{A}[t] = \text{const}_\sigma \mathbb{A}$  iff

$$\nexists g \in \mathbb{A} : \boxed{\sigma(g) = g + f}$$

## Represent sums (extension of Karr's result, 1981)

- ▶ Let  $(\mathbb{A}, \sigma)$  be a difference ring with constant field

$$\text{const}_\sigma \mathbb{A} := \{k \in \mathbb{A} \mid \sigma(k) = k\}.$$

- ▶ Adjoin a new variable  $t$  to  $\mathbb{A}$  (i.e.,  $\mathbb{A}[t]$  is a polynomial ring).
- ▶ Extend the shift operator s.t.

$$\sigma(t) = t + f \quad \text{for some } f \in \mathbb{A}.$$

Then  $\text{const}_\sigma \mathbb{A}[t] = \text{const}_\sigma \mathbb{A}$  iff

$$\nexists g \in \mathbb{A} : \boxed{\sigma(g) = g + f}$$

Such a difference ring extension  $(\mathbb{A}[t], \sigma)$  of  $(\mathbb{A}, \sigma)$  is called  $\Sigma^*$ -extension

## Represent sums (extension of Karr's result, 1981)

- ▶ Let  $(\mathbb{A}, \sigma)$  be a difference ring with constant field

$$\text{const}_\sigma \mathbb{A} := \{k \in \mathbb{A} \mid \sigma(k) = k\}.$$

- ▶ Adjoin a new variable  $t$  to  $\mathbb{A}$  (i.e.,  $\mathbb{A}[t]$  is a polynomial ring).
- ▶ Extend the shift operator s.t.

$$\sigma(t) = t + f \quad \text{for some } f \in \mathbb{A}.$$

Then  $\text{const}_\sigma \mathbb{A}[t] = \text{const}_\sigma \mathbb{A}$  iff

$$\nexists g \in \mathbb{A} : \boxed{\sigma(g) = g + f}$$

There are 2 cases:

1.  $\boxed{\nexists g \in \mathbb{A} : \sigma(g) = g + f}$ :  $(\mathbb{A}[t], \sigma)$  is a  $\Sigma^*$ -extension of  $(\mathbb{A}, \sigma)$

## Represent sums (extension of Karr's result, 1981)

- ▶ Let  $(\mathbb{A}, \sigma)$  be a difference ring with constant field

$$\text{const}_\sigma \mathbb{A} := \{k \in \mathbb{A} \mid \sigma(k) = k\}.$$

- ▶ Adjoin a new variable  $t$  to  $\mathbb{A}$  (i.e.,  $\mathbb{A}[t]$  is a polynomial ring).
- ▶ Extend the shift operator s.t.

$$\sigma(t) = t + f \quad \text{for some } f \in \mathbb{A}.$$

Then  $\text{const}_\sigma \mathbb{A}[t] = \text{const}_\sigma \mathbb{A}$  iff

$$\nexists g \in \mathbb{A} : \boxed{\sigma(g) = g + f}$$

There are 2 cases:

1.  $\boxed{\nexists g \in \mathbb{A} : \sigma(g) = g + f}$ :  $(\mathbb{A}[t], \sigma)$  is a  $\Sigma^*$ -extension of  $(\mathbb{A}, \sigma)$
2.  $\boxed{\exists g \in \mathbb{A} : \sigma(g) = g + f}$ : No need for a  $\Sigma^*$ -extension!

A hypergeometric *RHS-extension* of  $(\mathbb{K}(x), \sigma)$  is

- ▶ a ring

$$\mathbb{A} := \mathbb{K}(x)[p_1, p_1^{-1}][p_2, p_2^{-1}] \cdots [p_e, p_e^{-1}][z][s_1][s_2][s_3] \cdots$$

- ▶ with an automorphism where  $\sigma(c) = c$  for all  $c \in \mathbb{K}$  and where

$$\sigma(x) = x + 1$$

$$\begin{array}{l} \text{hypergeometric} \\ \text{products} \end{array} \quad \Leftrightarrow \quad \begin{array}{ll} \sigma(p_1) = a_1 p_1 & a_1 \in \mathbb{K}(x)^* \\ \sigma(p_2) = a_2 p_2 & a_2 \in \mathbb{K}(x)^* \end{array}$$

$$\vdots$$

$$\sigma(p_e) = a_e p_e \quad a_e \in \mathbb{K}(x)^*$$

$\gamma$  is a primitive  $\lambda$ th  
root of unity

 $\gamma^k$ 

$$\Leftrightarrow \quad \sigma(\mathbf{z}) = \gamma \mathbf{z} \quad \mathbf{z}^\lambda = \mathbf{1}$$

$$\begin{array}{l} \text{(nested) sum} \end{array} \quad \Leftrightarrow \quad \begin{array}{ll} \sigma(s_1) = s_1 + f_1 & f_1 \in \mathbb{K}(x)[p_1, p_1^{-1}] \cdots [p_e, p_e^{-1}][z] \\ \sigma(s_2) = s_2 + f_2 & f_2 \in \mathbb{K}(x)[p_1, p_1^{-1}] \cdots [p_e, p_e^{-1}][z][s_1] \\ \sigma(s_3) = s_3 + f_3 & f_3 \in \mathbb{K}(x)[p_1, p_1^{-1}] \cdots [p_e, p_e^{-1}][z][s_1][s_2] \\ \vdots & \end{array}$$

such that  $\text{const}_\sigma \mathbb{E} = \mathbb{K}$

## Represent products (extension of Karr's result, 1981)

- ▶ Let  $(\mathbb{A}, \sigma)$  be a difference ring with constant field

$$\text{const}_\sigma \mathbb{A} := \{k \in \mathbb{A} \mid \sigma(k) = k\}.$$

## Represent products (extension of Karr's result, 1981)

- ▶ Let  $(\mathbb{A}, \sigma)$  be a difference ring with constant field

$$\text{const}_\sigma \mathbb{A} := \{k \in \mathbb{A} \mid \sigma(k) = k\}.$$

- ▶ Take the ring of Laurent polynomials  $\mathbb{A}[t, \frac{1}{t}]$ .



## Represent products (extension of Karr's result, 1981)

- ▶ Let  $(\mathbb{A}, \sigma)$  be a difference ring with constant field

$$\text{const}_\sigma \mathbb{A} := \{k \in \mathbb{A} \mid \sigma(k) = k\}.$$

- ▶ Take the ring of Laurent polynomials  $\mathbb{A}[t, \frac{1}{t}]$ .
- ▶ Extend the shift operator s.t.

$$\sigma(t) = a t \quad \text{for some } a \in \mathbb{A}^*.$$

## Represent products (extension of Karr's result, 1981)

- ▶ Let  $(\mathbb{A}, \sigma)$  be a difference ring with constant field

$$\text{const}_\sigma \mathbb{A} := \{k \in \mathbb{A} \mid \sigma(k) = k\}.$$

- ▶ Take the ring of Laurent polynomials  $\mathbb{A}[t, \frac{1}{t}]$ .
- ▶ Extend the shift operator s.t.

$$\sigma(t) = at \quad \text{for some } a \in \mathbb{A}^*.$$

Then  $\text{const}_\sigma \mathbb{A}[t, t^{-1}] = \text{const}_\sigma \mathbb{A}$  iff

$$\nexists g \in \mathbb{A} \setminus \{0\} :$$

$$\boxed{\sigma(g) = ag}$$

## Represent products (extension of Karr's result, 1981)

- ▶ Let  $(\mathbb{A}, \sigma)$  be a difference ring with constant field

$$\text{const}_\sigma \mathbb{A} := \{k \in \mathbb{A} \mid \sigma(k) = k\}.$$

- ▶ Take the ring of Laurent polynomials  $\mathbb{A}[t, \frac{1}{t}]$ .
- ▶ Extend the shift operator s.t.

$$\sigma(t) = a t \quad \text{for some } a \in \mathbb{A}^*.$$

Then  $\text{const}_\sigma \mathbb{A}[t, t^{-1}] = \text{const}_\sigma \mathbb{A}$  iff



$\nexists g \in \mathbb{A} \setminus \{0\} :$

$$\sigma(g) = a g$$

## Represent products (extension of Karr's result, 1981)

- ▶ Let  $(\mathbb{A}, \sigma)$  be a difference ring with constant field

$$\text{const}_\sigma \mathbb{A} := \{k \in \mathbb{A} \mid \sigma(k) = k\}.$$

- ▶ Take the ring of Laurent polynomials  $\mathbb{A}[t, \frac{1}{t}]$ .
- ▶ Extend the shift operator s.t.

$$\sigma(t) = a t \quad \text{for some } a \in \mathbb{A}^*.$$

Then  $\text{const}_\sigma \mathbb{A}[t, t^{-1}] = \text{const}_\sigma \mathbb{A}$  iff

$$\nexists g \in \mathbb{A} \setminus \{0\} \nexists n \in \mathbb{Z} \setminus \{0\} : \boxed{\sigma(g) = a^n g}$$

## Represent products (extension of Karr's result, 1981)

- ▶ Let  $(\mathbb{A}, \sigma)$  be a difference ring with constant field

$$\text{const}_\sigma \mathbb{A} := \{k \in \mathbb{A} \mid \sigma(k) = k\}.$$

- ▶ Take the ring of Laurent polynomials  $\mathbb{A}[t, \frac{1}{t}]$ .
- ▶ Extend the shift operator s.t.

$$\sigma(t) = a t \quad \text{for some } a \in \mathbb{A}^*.$$

Then  $\text{const}_\sigma \mathbb{A}[t, t^{-1}] = \text{const}_\sigma \mathbb{A}$  iff

$$\nexists g \in \mathbb{A} \setminus \{0\} \nexists n \in \mathbb{Z} \setminus \{0\} : \boxed{\sigma(g) = a^n g}$$

Such a difference ring extension  $(\mathbb{A}[t, \frac{1}{t}], \sigma)$  of  $(\mathbb{A}, \sigma)$  is called **II-extension**

## Represent products (extension of Karr's result, 1981)

- ▶ Let  $(\mathbb{A}, \sigma)$  be a difference ring with constant field

$$\text{const}_\sigma \mathbb{A} := \{k \in \mathbb{A} \mid \sigma(k) = k\}.$$

- ▶ Take the ring of Laurent polynomials  $\mathbb{A}[t, \frac{1}{t}]$ .
- ▶ Extend the shift operator s.t.

$$\sigma(t) = at \quad \text{for some } a \in \mathbb{A}^*.$$

Then  $\text{const}_\sigma \mathbb{A}[t, t^{-1}] = \text{const}_\sigma \mathbb{A}$  iff

$$\nexists g \in \mathbb{A} \setminus \{0\} \nexists n \in \mathbb{Z} \setminus \{0\} : \boxed{\sigma(g) = a^n g}$$

There are 3 cases:

1.  $\boxed{\nexists g \in \mathbb{A} \setminus \{0\} \nexists n \in \mathbb{Z} \setminus \{0\} : \sigma(g) = a^n g}$ :  $(\mathbb{A}[t, \frac{1}{t}], \sigma)$  is a  $\Pi$ -ext. of  $(\mathbb{A}, \sigma)$

## Represent products (extension of Karr's result, 1981)

- ▶ Let  $(\mathbb{A}, \sigma)$  be a difference ring with constant field

$$\text{const}_\sigma \mathbb{A} := \{k \in \mathbb{A} \mid \sigma(k) = k\}.$$

- ▶ Take the ring of Laurent polynomials  $\mathbb{A}[t, \frac{1}{t}]$ .
- ▶ Extend the shift operator s.t.

$$\sigma(t) = at \quad \text{for some } a \in \mathbb{A}^*.$$

Then  $\text{const}_\sigma \mathbb{A}[t, t^{-1}] = \text{const}_\sigma \mathbb{A}$  iff

$$\nexists g \in \mathbb{A} \setminus \{0\} \nexists n \in \mathbb{Z} \setminus \{0\} : \boxed{\sigma(g) = a^n g}$$

There are 3 cases:

1.  $\boxed{\nexists g \in \mathbb{A} \setminus \{0\} \nexists n \in \mathbb{Z} \setminus \{0\} : \sigma(g) = a^n g}$ :  $(\mathbb{A}[t, \frac{1}{t}], \sigma)$  is a  $\Pi$ -ext. of  $(\mathbb{A}, \sigma)$
2.  $\boxed{\exists g \in \mathbb{A} \setminus \{0\} : \sigma(g) = ag}$ : No need for a  $\Pi$ -extension!

## Represent products (extension of Karr's result, 1981)

- ▶ Let  $(\mathbb{A}, \sigma)$  be a difference ring with constant field

$$\text{const}_\sigma \mathbb{A} := \{k \in \mathbb{A} \mid \sigma(k) = k\}.$$

- ▶ Take the ring of Laurent polynomials  $\mathbb{A}[t, \frac{1}{t}]$ .
- ▶ Extend the shift operator s.t.

$$\sigma(t) = at \quad \text{for some } a \in \mathbb{A}^*.$$

Then  $\text{const}_\sigma \mathbb{A}[t, t^{-1}] = \text{const}_\sigma \mathbb{A}$  iff

$$\nexists g \in \mathbb{A} \setminus \{0\} \nexists n \in \mathbb{Z} \setminus \{0\} : \boxed{\sigma(g) = a^n g}$$

There are 3 cases:

1.  $\boxed{\nexists g \in \mathbb{A} \setminus \{0\} \nexists n \in \mathbb{Z} \setminus \{0\} : \sigma(g) = a^n g}$ :  $(\mathbb{A}[t, \frac{1}{t}], \sigma)$  is a  $\Pi$ -ext. of  $(\mathbb{A}, \sigma)$
2.  $\boxed{\exists g \in \mathbb{A} \setminus \{0\} : \sigma(g) = ag}$ : No need for a  $\Pi$ -extension!
3.  $\boxed{\exists g \in \mathbb{A} \setminus \{0\} : \sigma(g) = a^n g \text{ only for } n \in \mathbb{Z} \setminus \{0, 1\}}$ : ☹️



## The hypergeometric case

- ▶ Take the difference field  $(\mathbb{K}(x), \sigma)$  with  $\sigma|_{\mathbb{K}} = \text{id}$  and  $\sigma(x) = x + 1$ .
- ▶ Let  $\alpha_1, \dots, \alpha_r \in \mathbb{K}(x)^*$

## The hypergeometric case

- ▶ Take the difference field  $(\mathbb{K}(x), \sigma)$  with  $\sigma|_{\mathbb{K}} = \text{id}$  and  $\sigma(x) = x + 1$ .
- ▶ Let  $\alpha_1, \dots, \alpha_r \in \mathbb{K}(x)^*$
- ▶ Then there is a difference ring

$$\mathbb{E}$$

such that for  $1 \leq i \leq r$  there are  $g_i \in \mathbb{E}^*$  with

$$\sigma(g_i) = \alpha_i g_i$$

## The hypergeometric case

- ▶ Take the difference field  $(\mathbb{K}(x), \sigma)$  with  $\sigma|_{\mathbb{K}} = \text{id}$  and  $\sigma(x) = x + 1$ .
- ▶ Let  $\alpha_1, \dots, \alpha_r \in \mathbb{K}(x)^*$
- ▶ Then there is a difference ring

$$\mathbb{E} = \mathbb{K}(x) \underbrace{[t_1, t_1^{-1}] \cdots [t_e, t_e^{-1}]}_{\text{tower of } \Pi\text{-ext.}} \underbrace{[z]}_{{(-1)^k \text{ or } \gamma^k}}$$

with

- ▶  $\frac{\sigma(t_i)}{t_i} \in \mathbb{K}(x)^*$  for  $1 \leq i \leq e$
- ▶  $\sigma(z) = \gamma z$  and  $z^\lambda = 1$  for some primitive  $\lambda$ th root of unity  $\gamma \in \mathbb{K}^*$
- ▶  $\text{const}_\sigma \mathbb{E} = \mathbb{K}$

such that for  $1 \leq i \leq r$  there are  $g_i \in \mathbb{E}^*$  with

$$\sigma(g_i) = \alpha_i g_i$$

## The hypergeometric case

- ▶ Take the difference field  $(\mathbb{K}(x), \sigma)$  with  $\sigma|_{\mathbb{K}} = \text{id}$  and  $\sigma(x) = x + 1$ .
- ▶ Let  $\alpha_1, \dots, \alpha_r \in \mathbb{K}(x)^*$
- ▶ Then there is a difference ring

$$\mathbb{E} = \mathbb{K}(x) \underbrace{[t_1, t_1^{-1}] \cdots [t_e, t_e^{-1}]}_{\text{tower of } \Pi\text{-ext.}} \underbrace{[z]}_{{(-1)^k \text{ or } \gamma^k}}$$

with

- ▶  $\frac{\sigma(t_i)}{t_i} \in \mathbb{K}(x)^*$  for  $1 \leq i \leq e$
- ▶  $\sigma(z) = \gamma z$  and  $z^\lambda = 1$  for some primitive  $\lambda$ th root of unity  $\gamma \in \mathbb{K}^*$
- ▶  $\text{const}_\sigma \mathbb{E} = \mathbb{K}$

such that for  $1 \leq i \leq r$  there are  $g_i \in \mathbb{E}^*$  with

$$\sigma(g_i) = \alpha_i g_i$$

Note: There are similar results for the  $q$ -rational, multi-basic and mixed case

A hypergeometric *RHS-extension* of  $(\mathbb{K}(x), \sigma)$  is

- ▶ a ring

$$\mathbb{A} := \mathbb{K}(x)[p_1, p_1^{-1}][p_2, p_2^{-1}] \cdots [p_e, p_e^{-1}][z][s_1][s_2][s_3] \cdots$$

- ▶ with an automorphism where  $\sigma(c) = c$  for all  $c \in \mathbb{K}$  and where

$$\sigma(x) = x + 1$$

$$\begin{array}{l} \text{hypergeometric} \\ \text{products} \end{array} \quad \Leftrightarrow \quad \begin{array}{ll} \sigma(p_1) = a_1 p_1 & a_1 \in \mathbb{K}(x)^* \\ \sigma(p_2) = a_2 p_2 & a_2 \in \mathbb{K}(x)^* \end{array}$$

$$\vdots$$

$$\sigma(p_e) = a_e p_e \quad a_e \in \mathbb{K}(x)^*$$

$\gamma$  is a primitive  $\lambda$ th  
root of unity

 $\gamma^k$ 

$$\Leftrightarrow \quad \sigma(\mathbf{z}) = \gamma \mathbf{z} \quad \mathbf{z}^\lambda = \mathbf{1}$$

$$\begin{array}{l} \text{(nested) sum} \end{array} \quad \Leftrightarrow \quad \begin{array}{ll} \sigma(s_1) = s_1 + f_1 & f_1 \in \mathbb{K}(x)[p_1, p_1^{-1}] \cdots [p_e, p_e^{-1}][z] \\ \sigma(s_2) = s_2 + f_2 & f_2 \in \mathbb{K}(x)[p_1, p_1^{-1}] \cdots [p_e, p_e^{-1}][z][s_1] \\ \sigma(s_3) = s_3 + f_3 & f_3 \in \mathbb{K}(x)[p_1, p_1^{-1}] \cdots [p_e, p_e^{-1}][z][s_1][s_2] \\ \vdots & \end{array}$$

such that  $\text{const}_\sigma \mathbb{E} = \mathbb{K}$

This yields a strategy (actually the only strategy for shift-stable sets).

## A Strategy to solve Problem SigmaReduce

Given:  $A_1, \dots, A_u \in \text{SumProd}(\mathbb{G})$  with  $\mathbb{G} = \mathbb{K}(x)$ .

Find: a  $\sigma$ -reduced set  $W = \{T_1, \dots, T_e\} \subset \Sigma\Pi(\mathbb{G})$  with  $B_1, \dots, B_u \in \text{SumProd}(W, \mathbb{G})$  and  $\delta_1, \dots, \delta_u \in \mathbb{N}$  such that  $A_i(n) = B_i(n)$  holds for all  $n \geq \delta_i$  and  $1 \leq i \leq r$ .

1. Construct  $R\Pi\Sigma$ -extension  $(\mathbb{E}, \sigma)$  of  $(\mathbb{G}, \sigma)$  with  $\mathbb{E} = \mathbb{G}\langle t_1 \rangle \dots \langle t_e \rangle$  equipped with  $\text{ev} : \mathbb{E} \times \mathbb{N} \rightarrow \mathbb{K}$  such that we get  $a_1, \dots, a_u \in \mathbb{E}$  and  $\delta_1, \dots, \delta_u \in \mathbb{N}$  with

$$A_i(n) = \text{ev}(a_i, n) \quad \forall n \geq \delta_i. \quad (12)$$

2. Set  $W = \{T_1, \dots, T_e\}$  with  $T_i := \text{expr}(t_i) \in \Sigma\Pi(\mathbb{G})$  for  $1 \leq i \leq e$ .
3. Set  $B_i := \text{expr}(a_i) \in \text{SumProd}(W, \mathbb{G})$  for  $1 \leq i \leq u$ .
4. Return  $W, (B_1, \dots, B_u)$  and  $(\delta_1, \dots, \delta_u)$ .

This yields a strategy (actually the only strategy for shift-stable sets).

### An Algorithm to solve Problem SigmaReduce

Given:  $A_1, \dots, A_u \in \text{SumProd}(\mathbb{G})$  with  $\mathbb{G} = \mathbb{K}(x)$ .

Find: a  $\sigma$ -reduced set  $W = \{T_1, \dots, T_e\} \subset \Sigma\Pi(\mathbb{G})$  with  $B_1, \dots, B_u \in \text{SumProd}(W, \mathbb{G})$  and  $\delta_1, \dots, \delta_u \in \mathbb{N}$  such that  $A_i(n) = B_i(n)$  holds for all  $n \geq \delta_i$  and  $1 \leq i \leq r$ .

1. Construct  $R\Pi\Sigma$ -extension  $(\mathbb{E}, \sigma)$  of  $(\mathbb{G}, \sigma)$  with  $\mathbb{E} = \mathbb{G}\langle t_1 \rangle \dots \langle t_e \rangle$  equipped with  $\text{ev} : \mathbb{E} \times \mathbb{N} \rightarrow \mathbb{K}$  such that we get  $a_1, \dots, a_u \in \mathbb{E}$  and  $\delta_1, \dots, \delta_u \in \mathbb{N}$  with

$$A_i(n) = \text{ev}(a_i, n) \quad \forall n \geq \delta_i. \quad (12)$$

2. Set  $W = \{T_1, \dots, T_e\}$  with  $T_i := \text{expr}(t_i) \in \Sigma\Pi(\mathbb{G})$  for  $1 \leq i \leq e$ .
3. Set  $B_i := \text{expr}(a_i) \in \text{SumProd}(W, \mathbb{G})$  for  $1 \leq i \leq u$ .
4. Return  $W, (B_1, \dots, B_u)$  and  $(\delta_1, \dots, \delta_u)$ .

## Telescoping

GIVEN  $f(k) = S_1(k)$ .

FIND  $g(k)$ :

$$f(k) = g(k+1) - g(k)$$

for all  $1 \leq k \leq n$  and  $n \geq 0$ .



## Telescoping

GIVEN  $f(k) = S_1(k)$ .

FIND  $g(k)$ :

$$f(k) = g(k+1) - g(k)$$

for all  $1 \leq k \leq n$  and  $n \geq 0$ .

Sigma computes

$$g(k) = (S_1(k) - 1)k.$$

## Telescoping

GIVEN  $f(k) = S_1(k)$ .

FIND  $g(k)$ :

$$f(k) = g(k + 1) - g(k)$$

for all  $1 \leq k \leq n$  and  $n \geq 0$ .

Summing this equation over  $k$  from 1 to  $n$  gives

$$\begin{aligned} \sum_{k=1}^n S_1(k) &= g(n + 1) - g(1) \\ &= (S_1(n + 1) - 1)(n + 1). \end{aligned}$$

## Telescoping in the given difference ring

FIND a closed form for

$$\sum_{k=1}^n S_1(k).$$

**A difference ring for the [summand](#)**

Consider a ring

$$\mathbb{A}$$

with the automorphism  $\sigma : \mathbb{A} \rightarrow \mathbb{A}$  defined by

## Telescoping in the given difference ring

FIND a closed form for

$$\sum_{k=1}^n S_1(k).$$

### A difference ring for the **summand**

Consider a ring

$$\mathbb{A} := \mathbb{Q}$$

with the automorphism  $\sigma : \mathbb{A} \rightarrow \mathbb{A}$  defined by

$$\sigma(c) = c \quad \forall c \in \mathbb{Q},$$

## Telescoping in the given difference ring

FIND a closed form for

$$\sum_{k=1}^n S_1(k).$$

### A difference ring for the **summand**

Consider a ring

$$\mathbb{A} := \mathbb{Q}(x)$$

with the automorphism  $\sigma : \mathbb{A} \rightarrow \mathbb{A}$  defined by

$$\sigma(c) = c \quad \forall c \in \mathbb{Q},$$

$$\sigma(x) = x + 1,$$

$$S k = k + 1,$$

## Telescoping in the given difference ring

FIND a closed form for

$$\sum_{k=1}^n S_1(k).$$

### A difference ring for the **summand**

Consider a ring

$$\mathbb{A} := \mathbb{Q}(x)[h]$$

with the automorphism  $\sigma : \mathbb{A} \rightarrow \mathbb{A}$  defined by

$$\sigma(c) = c \quad \forall c \in \mathbb{Q},$$

$$\sigma(x) = x + 1,$$

$$\sigma(h) = h + \frac{1}{x+1},$$

$$\mathcal{S}k = k + 1,$$

$$\mathcal{S}S_1(k) = S_1(k) + \frac{1}{k+1}.$$

## Telescoping in the given difference ring

FIND  $g \in \mathbb{A}$ :

$$\sigma(g) - g = h.$$

## Telescoping in the given difference ring

FIND  $g \in \mathbb{A}$ :

$$\sigma(g) - g = h.$$

We compute

$$g = (h - 1)x \in \mathbb{A}.$$



## Telescoping in the given difference ring

FIND  $g \in \mathbb{A}$ :

$$\sigma(g) - g = h.$$

We compute

$$g = (h - 1)x \in \mathbb{A}.$$

This gives

$$g(k+1) - g(k) = S_1(k)$$

with

$$g(k) = (S_1(k) - 1)k.$$

## Telescoping in the given difference ring

FIND  $g \in \mathbb{A}$ :

$$\sigma(g) - g = h.$$

We compute

$$g = (h - 1)x \in \mathbb{A}.$$

This gives

$$g(k+1) - g(k) = S_1(k)$$

with

$$g(k) = (S_1(k) - 1)k.$$

Hence,

$$(S_1(n+1) - 1)(n+1) = \sum_{k=1}^n S_1(k).$$

FIND  $g \in \mathbb{Q}(x)[h]$ :

$$\sigma(g) - g = h.$$

FIND  $g \in \mathbb{Q}(x)[h]$ :

$$\sigma(g) - g = h.$$

**Degree bound:** COMPUTE  $b \geq 0$ :

$$\forall g \in \mathbb{Q}(x)[h] \quad \sigma(g) - g = h \quad \Rightarrow \quad \deg(g) \leq b.$$

FIND  $g \in \mathbb{Q}(x)[h]$ :

$$\sigma(g) - g = h.$$

**Degree bound:** COMPUTE  $b \geq 0$ :

$$b = 2$$

$$\forall g \in \mathbb{Q}(x)[h] \quad \sigma(g) - g = h \quad \Rightarrow \quad \deg(g) \leq b.$$

FIND  $g \in \mathbb{Q}(x)[h]$ :

$$\sigma(g) - g = h.$$

**Degree bound:** COMPUTE  $b \geq 0$ :

$$\forall g \in \mathbb{Q}(x)[h] \quad \sigma(g) - g = h \quad \Rightarrow \quad \deg(g) \leq b.$$

**Polynomial Solution:** FIND

$$g = g_2 h^2 + g_1 h + g_0 \in \mathbb{Q}(x)[h].$$

ANSATZ  $g = g_2 h^2 + g_1 h + g_0 \in \mathbb{Q}(x)[h]$

$$\sigma(g) - g = h$$



ANSATZ  $g = g_2 h^2 + g_1 h + g_0 \in \mathbb{Q}(x)[h]$

$$\begin{aligned} & [\sigma(g_2 h^2 + g_1 h + g_0)] \\ & - [g_2 h^2 + g_1 h + g_0] = h \end{aligned}$$





ANSATZ  $g = g_2 h^2 + g_1 h + g_0 \in \mathbb{Q}(x)[h]$

$$\begin{aligned} & [\sigma(g_2 h^2) + \sigma(g_1 h + g_0)] \\ & \quad - [g_2 h^2 + g_1 h + g_0] = h \end{aligned}$$



ANSATZ  $g = g_2 h^2 + g_1 h + g_0 \in \mathbb{Q}(x)[h]$

$$\begin{aligned} & [\sigma(g_2) \sigma(h^2) + \sigma(g_1 h + g_0)] \\ & \quad - [g_2 h^2 + g_1 h + g_0] = h \end{aligned}$$



ANSATZ  $g = g_2 h^2 + g_1 h + g_0 \in \mathbb{Q}(x)[h]$

$$\begin{aligned} & [\sigma(g_2) \sigma(h)^2 + \sigma(g_1 h + g_0)] \\ & \quad - [g_2 h^2 + g_1 h + g_0] = h \end{aligned}$$



ANSATZ  $g = g_2 h^2 + g_1 h + g_0 \in \mathbb{Q}(x)[h]$

$$\begin{aligned} & \left[ \sigma(g_2) \left( h + \frac{1}{x+1} \right)^2 + \sigma(g_1 h + g_0) \right] \\ & \quad - [g_2 h^2 + g_1 h + g_0] = h \end{aligned}$$



$$\text{ANSATZ } g = g_2 h^2 + g_1 h + g_0 \in \mathbb{Q}(x)[h]$$

$$\left[ \sigma(g_2) \left( h + \frac{1}{x+1} \right)^2 + \sigma(g_1 h + g_0) \right] - [g_2 h^2 + g_1 h + g_0] = h$$

coeff. comp. 

$$\sigma(g_2) - g_2 = 0$$

$$\text{ANSATZ } g = g_2 h^2 + g_1 h + g_0 \in \mathbb{Q}(x)[h]$$

$$\left[ \sigma(g_2) \left( h + \frac{1}{x+1} \right)^2 + \sigma(g_1 h + g_0) \right] - [g_2 h^2 + g_1 h + g_0] = h$$

coeff. comp. 

$$\sigma(g_2) - g_2 = 0$$

$$g_2 = c \in \mathbb{Q}$$

ANSATZ  $g = g_2 h^2 + g_1 h + g_0 \in \mathbb{Q}(x)[h]$

$$\left[ \sigma(g_2) \left( h + \frac{1}{x+1} \right)^2 + \sigma(g_1 h + g_0) \right] - [g_2 h^2 + g_1 h + g_0] = h$$

coeff. comp.

$$\sigma(g_2) - g_2 = 0$$

$$g_2 = c \in \mathbb{Q}$$

$$\left[ \sigma(c) \left( h + \frac{1}{x+1} \right)^2 + \sigma(g_1 h + g_0) \right] - [c h^2 + g_1 h + g_0] = h$$

$$\text{ANSATZ } g = g_2 h^2 + g_1 h + g_0 \in \mathbb{Q}(x)[h]$$

$$\left[ \sigma(g_2) \left( h + \frac{1}{x+1} \right)^2 + \sigma(g_1 h + g_0) \right] - [g_2 h^2 + g_1 h + g_0] = h$$

coeff. comp.

$$\sigma(g_2) - g_2 = 0$$

$$g_2 = c \in \mathbb{Q}$$

$$\left[ c \left( h + \frac{1}{x+1} \right)^2 + \sigma(g_1 h + g_0) \right] - [c h^2 + g_1 h + g_0] = h$$



$$\text{ANSATZ } g = g_2 h^2 + g_1 h + g_0 \in \mathbb{Q}(x)[h]$$

$$\begin{aligned} & [\sigma(g_2)(h + \frac{1}{x+1})^2 + \sigma(g_1 h + g_0)] \\ & - [g_2 h^2 + g_1 h + g_0] = h \end{aligned}$$

coeff. comp. 

$$\sigma(g_2) - g_2 = 0$$

$$g_2 = c \in \mathbb{Q}$$

$$\sigma(g_1 h + g_0) - (g_1 h + g_0) = h - c \left[ \frac{2h(x+1)+1}{(x+1)^2} \right]$$

$$\text{ANSATZ } g = g_2 h^2 + g_1 h + g_0 \in \mathbb{Q}(x)[h]$$

$$[\sigma(g_2)(h + \frac{1}{x+1})^2 + \sigma(g_1 h + g_0)] - [g_2 h^2 + g_1 h + g_0] = h$$

coeff. comp.

$$\sigma(g_2) - g_2 = 0$$

$$g_2 = c \in \mathbb{Q}$$

$$\sigma(g_1 h + g_0) - (g_1 h + g_0) = h - c \left[ \frac{2h(x+1)+1}{(x+1)^2} \right]$$

coeff. comp.

$$\sigma(g_1) - g_1 = 1 - c \frac{2}{x+1}$$

$$\text{ANSATZ } g = g_2 h^2 + g_1 h + g_0 \in \mathbb{Q}(x)[h]$$

$$\left[ \sigma(g_2) \left( h + \frac{1}{x+1} \right)^2 + \sigma(g_1 h + g_0) \right] - [g_2 h^2 + g_1 h + g_0] = h$$

coeff. comp.

$$\sigma(g_2) - g_2 = 0$$

$$g_2 = c \in \mathbb{Q}$$

$$\sigma(g_1 h + g_0) - (g_1 h + g_0) = h - c \left[ \frac{2h(x+1)+1}{(x+1)^2} \right]$$

coeff. comp.

$$\sigma(g_1) - g_1 = 1 - c \frac{2}{x+1}$$

$$c = 0, \quad g_1 = x + d \\ d \in \mathbb{Q}$$

$$\text{ANSATZ } g = g_2 h^2 + g_1 h + g_0 \in \mathbb{Q}(x)[h]$$

$$\left[ \sigma(g_2) \left( h + \frac{1}{x+1} \right)^2 + \sigma(g_1 h + g_0) \right] - [g_2 h^2 + g_1 h + g_0] = h$$

coeff. comp. 

$$\sigma(g_2) - g_2 = 0$$

$$g_2 = c \in \mathbb{Q}$$

$$\sigma(g_1 h + g_0) - (g_1 h + g_0) = h - c \left[ \frac{2h(x+1)+1}{(x+1)^2} \right]$$

coeff. comp.

$$\sigma(g_1) - g_1 = 1 - c \frac{2}{x+1}$$

$$\sigma(g_0) - g_0 = -1 - d \frac{1}{x+1}$$

$$c = 0, \quad g_1 = x + d \\ d \in \mathbb{Q}$$

$$\text{ANSATZ } g = g_2 h^2 + g_1 h + g_0 \in \mathbb{Q}(x)[h]$$

$$\left[ \sigma(g_2) \left( h + \frac{1}{x+1} \right)^2 + \sigma(g_1 h + g_0) \right] - [g_2 h^2 + g_1 h + g_0] = h$$

coeff. comp.

$$g = hx - x$$

$$\sigma(g_2) - g_2 = 0$$

$$g_2 = c \in \mathbb{Q}$$

$$\sigma(g_1 h + g_0) - (g_1 h + g_0) = h - c \left[ \frac{2h(x+1)+1}{(x+1)^2} \right]$$

coeff. comp.

$$\sigma(g_1) - g_1 = 1 - c \frac{2}{x+1}$$

$$\begin{aligned} g_0 &= -x \\ d &= 0 \end{aligned}$$

$$\left\langle \sigma(g_0) - g_0 = -1 - d \frac{1}{x+1} \right.$$

$$c = 0, \quad g_1 = x + d \\ d \in \mathbb{Q}$$

## Telescoping in the given difference ring

FIND  $g \in \mathbb{A}$ :

$$\sigma(g) - g = h.$$

We compute

$$g = (h - 1)x \in \mathbb{A}.$$

This gives

$$g(k+1) - g(k) = S_1(k)$$

with

$$g(k) = (S_1(k) - 1)k.$$

Hence,

$$(S_1(n+1) - 1)(n+1) = \sum_{k=1}^n S_1(k).$$

**Remarks.** All results can be generalized to the following setting:

- ▶ the **mixed multibasic hypergeometric case**:

$\mathbb{G} := \mathbb{K}(x, x_1, \dots, x_v)$  with  $\mathbb{K} = K(q_1, \dots, q_v)$  For  $f = \frac{p}{q} \in \mathbb{G}$  with  $p, q \in \mathbb{K}[x, x_1, \dots, x_v]$  where  $q \neq 0$  and  $p, q$  being coprime we define

$$\text{ev}(f, k) = \begin{cases} 0 & \text{if } q(k, q_1^k, \dots, q_v^k) = 0 \\ \frac{p(k, q_1^k, \dots, q_v^k)}{q(k, q_1^k, \dots, q_v^k)} & \text{if } q(k, q_1^k, \dots, q_v^k) \neq 0. \end{cases}$$

**Remarks.** All results can be generalized to the following setting:

- ▶ the **mixed multibasic hypergeometric case**:

$\mathbb{G} := \mathbb{K}(x, x_1, \dots, x_v)$  with  $\mathbb{K} = K(q_1, \dots, q_v)$  For  $f = \frac{p}{q} \in \mathbb{G}$  with  $p, q \in \mathbb{K}[x, x_1, \dots, x_v]$  where  $q \neq 0$  and  $p, q$  being coprime we define

$$\text{ev}(f, k) = \begin{cases} 0 & \text{if } q(k, q_1^k, \dots, q_v^k) = 0 \\ \frac{p(k, q_1^k, \dots, q_v^k)}{q(k, q_1^k, \dots, q_v^k)} & \text{if } q(k, q_1^k, \dots, q_v^k) \neq 0. \end{cases}$$

- ▶ simple products:  $\text{Prod}^*(\mathbb{G})$  is the smallest set that contains 1 with:

1. If  $r \in \mathcal{R}$  then  $\text{RPow}(r) \in \text{Prod}^*(\mathbb{G})$ .
2. If  $f \in \mathbb{G}^*$ ,  $l \in \mathbb{N}$  with  $l \geq Z(f)$  then  $\text{Prod}(l, f) \in \text{Prod}^*(\mathbb{G})$ .
3. If  $p, q \in \text{Prod}^*(\mathbb{G})$  then  $p \odot q \in \text{Prod}^*(\mathbb{G})$ .
4. If  $p \in \text{Prod}^*(\mathbb{G})$  and  $z \in \mathbb{Z} \setminus \{0\}$  then  $p^{\otimes z} \in \text{Prod}^*(\mathbb{G})$ .



**Remarks.** All results can be generalized to the following setting:

- ▶ the **mixed multibasic hypergeometric case**:

$\mathbb{G} := \mathbb{K}(x, x_1, \dots, x_v)$  with  $\mathbb{K} = K(q_1, \dots, q_v)$  For  $f = \frac{p}{q} \in \mathbb{G}$  with  $p, q \in \mathbb{K}[x, x_1, \dots, x_v]$  where  $q \neq 0$  and  $p, q$  being coprime we define

$$\text{ev}(f, k) = \begin{cases} 0 & \text{if } q(k, q_1^k, \dots, q_v^k) = 0 \\ \frac{p(k, q_1^k, \dots, q_v^k)}{q(k, q_1^k, \dots, q_v^k)} & \text{if } q(k, q_1^k, \dots, q_v^k) \neq 0. \end{cases}$$

- ▶ **nested products**:  $\text{Prod}^*(\mathbb{G})$  is the smallest set that contains 1 with:
  1. If  $r \in \mathcal{R}$  then  $\text{RPow}(r) \in \text{Prod}^*(\mathbb{G})$ .
  2. If  $p \in \text{Prod}^*(\mathbb{G})$ ,  $f \in \mathbb{G}^*$ ,  $l \in \mathbb{N}$  with  $l \geq Z(f)$  then  $\text{Prod}(l, f \odot p) \in \text{Prod}^*(\mathbb{G})$ .
  3. If  $p, q \in \text{Prod}^*(\mathbb{G})$  then  $p \odot q \in \text{Prod}^*(\mathbb{G})$ .
  4. If  $p \in \text{Prod}^*(\mathbb{G})$  and  $z \in \mathbb{Z} \setminus \{0\}$  then  $p^{\otimes z} \in \text{Prod}^*(\mathbb{G})$ .

**Remarks.** All results can be generalized to the following setting:

- ▶ the **mixed multibasic hypergeometric case**:

$\mathbb{G} := \mathbb{K}(x, x_1, \dots, x_v)$  with  $\mathbb{K} = K(q_1, \dots, q_v)$  For  $f = \frac{p}{q} \in \mathbb{G}$  with  $p, q \in \mathbb{K}[x, x_1, \dots, x_v]$  where  $q \neq 0$  and  $p, q$  being coprime we define

$$\text{ev}(f, k) = \begin{cases} 0 & \text{if } q(k, q_1^k, \dots, q_v^k) = 0 \\ \frac{p(k, q_1^k, \dots, q_v^k)}{q(k, q_1^k, \dots, q_v^k)} & \text{if } q(k, q_1^k, \dots, q_v^k) \neq 0. \end{cases}$$

- ▶ **nested products**:  $\text{Prod}^*(\mathbb{G})$  is the smallest set that contains 1 with:
  1. If  $r \in \mathcal{R}$  then  $\text{RPow}(r) \in \text{Prod}^*(\mathbb{G})$ .
  2. If  $p \in \text{Prod}^*(\mathbb{G})$ ,  $f \in \mathbb{G}^*$ ,  $l \in \mathbb{N}$  with  $l \geq Z(f)$  then  $\text{Prod}(l, f \odot p) \in \text{Prod}^*(\mathbb{G})$ .
  3. If  $p, q \in \text{Prod}^*(\mathbb{G})$  then  $p \odot q \in \text{Prod}^*(\mathbb{G})$ .
  4. If  $p \in \text{Prod}^*(\mathbb{G})$  and  $z \in \mathbb{Z} \setminus \{0\}$  then  $p^{\otimes z} \in \text{Prod}^*(\mathbb{G})$ .

For further details see

Term Algebras, Canonical Representations and Difference Ring Theory for Symbolic Summation. To appear in: Anti-Differentiation and the Calculation of Feynman Amplitudes, J. Blümlein and C. Schneider (ed.), Texts and Monographs in Symbolic Computation, 2021. Springer, arXiv:2102.01471 [cs.SC]

## General picture:

Part 1: Symbolic summation (a short introduction)

Part 2: Modeling of sequences with a term algebra (user interface)

Part 3: Modeling of sequences in difference rings (computer algebra)

Part 4: Construction of appropriate difference rings (advanced CA)

Part 5: Applications

## A warm-up example: simplify

$$\sum_{k=0}^{\infty} \sum_{j=0}^{\infty} \left( \frac{(2j+k+n+2)j!k!(j+k+n)!}{(j+k+1)(j+n+1)(j+k+1)!(j+n+1)!(k+n+1)!} \right. \\ \left. + \underbrace{\frac{j!k!(j+k+n)!(-S_1(j) + S_1(j+k) + S_1(j+n) - S_1(j+k+n))}{(j+k+1)!(j+n+1)!(k+n+1)!}}_{f(j)} \right)$$

where

$$S_1(n) = \sum_{i=1}^n \frac{1}{i} \quad (= H_n)$$

Arose in the context of

I. Bierenbaum, J. Blümlein, and S. Klein, **Evaluating two-loop massive operator matrix elements with Mellin-Barnes integrals.** 2006

## A warm-up example: simplify

$$\sum_{k=0}^{\infty} \sum_{j=0}^{\infty} \left( \frac{(2j+k+n+2)j!k!(j+k+n)!}{(j+k+1)(j+n+1)(j+k+1)!(j+n+1)!(k+n+1)!} \right. \\ \left. + \underbrace{\frac{j!k!(j+k+n)!(-S_1(j) + S_1(j+k) + S_1(j+n) - S_1(j+k+n))}{(j+k+1)!(j+n+1)!(k+n+1)!}}_{f(j)} \right)$$

FIND  $g(j)$ :

$$\boxed{f(j) = g(j+1) - g(j)}$$

## A warm-up example: simplify

$$\sum_{k=0}^{\infty} \sum_{j=0}^{\infty} \left( \frac{(2j+k+n+2)j!k!(j+k+n)!}{(j+k+1)(j+n+1)(j+k+1)!(j+n+1)!(k+n+1)!} \right. \\ \left. + \underbrace{\frac{j!k!(j+k+n)!(-S_1(j) + S_1(j+k) + S_1(j+n) - S_1(j+k+n))}{(j+k+1)!(j+n+1)!(k+n+1)!}}_{f(j)} \right)$$

FIND  $g(j)$ :

$$\boxed{f(j) = g(j+1) - g(j)}$$

↑ summation package Sigma

$$g(j) = \frac{(j+k+1)(j+n+1)j!k!(j+k+n)!(S_1(j) - S_1(j+k) - S_1(j+n) + S_1(j+k+n))}{kn(j+k+1)!(j+n+1)!(k+n+1)!}$$

## A warm-up example: simplify

$$\sum_{k=0}^{\infty} \sum_{j=0}^{\infty} \left( \frac{(2j+k+n+2)j!k!(j+k+n)!}{(j+k+1)(j+n+1)(j+k+1)!(j+n+1)!(k+n+1)!} \right. \\ \left. + \underbrace{\frac{j!k!(j+k+n)!(-S_1(j) + S_1(j+k) + S_1(j+n) - S_1(j+k+n))}{(j+k+1)!(j+n+1)!(k+n+1)!}}_{f(j)} \right)$$

FIND  $g(j)$ :

$$\boxed{f(j) = g(j+1) - g(j)}$$

Summing the telescoping equation over  $j$  from 0 to  $a$  gives

$$\sum_{j=0}^a f(j) = g(a+1) - g(0)$$

A warm-up example: simplify

$$\sum_{k=0}^{\infty} \sum_{j=0}^{\infty} \left( \frac{(2j+k+n+2)j!k!(j+k+n)!}{(j+k+1)(j+n+1)(j+k+1)!(j+n+1)!(k+n+1)!} \right. \\ \left. + \frac{j!k!(j+k+n)!(-S_1(j) + S_1(j+k) + S_1(j+n) - S_1(j+k+n))}{(j+k+1)!(j+n+1)!(k+n+1)!} \right) \\ \underbrace{\hspace{15em}}_{f(j)}$$

FIND  $g(j)$ :

$$\boxed{f(j) = g(j+1) - g(j)}$$

Summing the telescoping equation over  $j$  from 0 to  $a$  gives

$$\sum_{j=0}^a f(j) = g(a+1) - g(0) \\ = \frac{(a+1)!(k-1)!(a+k+n+1)!(S_1(a) - S_1(a+k) - S_1(a+n) + S_1(a+k+n))}{n(a+k+1)!(a+n+1)!(k+n+1)!} \\ + \underbrace{\frac{S_1(k) + S_1(n) - S_1(k+n)}{kn(k+n+1)n!} + \frac{(2a+k+n+2)a!k!(a+k+n)!}{(a+k+1)(a+n+1)(a+k+1)!(a+n+1)!(k+n+1)!}}_{a \rightarrow \infty}$$



## A warm-up example: simplify

$$\sum_{k=0}^{\infty} \sum_{j=0}^{\infty} \left( \frac{(2j+k+n+2)j!k!(j+k+n)!}{(j+k+1)(j+n+1)(j+k+1)!(j+n+1)!(k+n+1)!} \right. \\ \left. + \frac{j!k!(j+k+n)!(-S_1(j) + S_1(j+k) + S_1(j+n) - S_1(j+k+n))}{(j+k+1)!(j+n+1)!(k+n+1)!} \right) \\ \underbrace{\hspace{15em}}_{f(j)}$$

$$\sum_{j=0}^{\infty} f(j) = \frac{S_1(k) + S_1(n) - S_1(k+n)}{kn(k+n+1)}$$

In[8]:= << Sigma.m

Sigma - A summation package by Carsten Schneider © RISC-Linz

$$\text{In[9]:= mySum} = \sum_{j=0}^a \left( \frac{(2j+k+n+2)j!k!(j+k+n)!}{(j+k+1)(j+n+1)(j+k+1)!(j+n+1)!(k+n+1)!} + \frac{j!k!(j+k+n)!(-S_1[j] + S_1[j+k] + S_1[j+n] - S_1[j+k+n])}{(j+k+1)!(j+n+1)!(k+n+1)!} \right);$$

In[8]:= &lt;&lt; Sigma.m

Sigma - A summation package by Carsten Schneider © RISC-Linz

$$\text{In[9]:= mySum} = \sum_{j=0}^a \left( \frac{(2j+k+n+2)j!k!(j+k+n)!}{(j+k+1)(j+n+1)(j+k+1)!(j+n+1)!(k+n+1)!} + \frac{j!k!(j+k+n)!(-S_1[j] + S_1[j+k] + S_1[j+n] - S_1[j+k+n])}{(j+k+1)!(j+n+1)!(k+n+1)!} \right);$$

In[10]:= res = SigmaReduce[mySum]

$$\text{Out[10]=} \frac{(a+1)!(k-1)!(a+k+n+1)!(S_1[a] - S_1[a+k] - S_1[a+n] + S_1[a+k+n])}{n(a+k+1)!(a+n+1)!(k+n+1)!} + \frac{S_1[k] + S_1[n] - S_1[k+n]}{kn(k+n+1)n!} + \frac{(2a+k+n+2)a!k!(a+k+n)!}{(a+k+1)(a+n+1)(a+k+1)!(a+n+1)!(k+n+1)!}$$

In[8]:= << Sigma.m

Sigma - A summation package by Carsten Schneider © RISC-Linz

$$\text{In[9]:= mySum} = \sum_{j=0}^a \left( \frac{(2j+k+n+2)j!k!(j+k+n)!}{(j+k+1)(j+n+1)(j+k+1)!(j+n+1)!(k+n+1)!} + \frac{j!k!(j+k+n)!(-S_1[j] + S_1[j+k] + S_1[j+n] - S_1[j+k+n])}{(j+k+1)!(j+n+1)!(k+n+1)!} \right);$$

In[10]:= res = SigmaReduce[mySum]

$$\text{Out[10]=} \frac{(a+1)!(k-1)!(a+k+n+1)!(S_1[a] - S_1[a+k] - S_1[a+n] + S_1[a+k+n])}{n(a+k+1)!(a+n+1)!(k+n+1)!} + \frac{S_1[k] + S_1[n] - S_1[k+n]}{kn(k+n+1)n!} + \frac{(2a+k+n+2)a!k!(a+k+n)!}{(a+k+1)(a+n+1)(a+k+1)!(a+n+1)!(k+n+1)!}$$

In[11]:= SigmaLimit[res, {n}, a]

$$\text{Out[11]=} \frac{1}{n!} \frac{S_1[k] + S_1[n] - S_1[k+n]}{kn(k+n+1)}$$

## A warm-up example: simplify

$$\sum_{k=0}^{\infty} \sum_{j=0}^{\infty} \left( \frac{(2j+k+n+2)j!k!(j+k+n)!}{(j+k+1)(j+n+1)(j+k+1)!(j+n+1)!(k+n+1)!} \right. \\ \left. + \frac{j!k!(j+k+n)!(-S_1(j) + S_1(j+k) + S_1(j+n) - S_1(j+k+n))}{(j+k+1)!(j+n+1)!(k+n+1)!} \right) \\ \underbrace{\hspace{15em}}_{f(j)}$$

$$\sum_{j=0}^{\infty} f(j) = \frac{S_1(k) + S_1(n) - S_1(k+n)}{kn(k+n+1)}$$

## A warm-up example: simplify

$$\sum_{k=0}^{\infty} \sum_{j=0}^{\infty} \left( \frac{(2j+k+n+2)j!k!(j+k+n)!}{(j+k+1)(j+n+1)(j+k+1)!(j+n+1)!(k+n+1)!} \right. \\ \left. + \frac{j!k!(j+k+n)!(-S_1(j) + S_1(j+k) + S_1(j+n) - S_1(j+k+n))}{(j+k+1)!(j+n+1)!(k+n+1)!} \right) \\ \underbrace{\hspace{15em}}_{f(j)}$$

$$\sum_{k=1}^{\infty} \sum_{j=0}^{\infty} f(j) = \frac{1}{n!} \sum_{k=1}^{\infty} \frac{S_1(k) + S_1(n) - S_1(k+n)}{kn(k+n+1)}$$

## Telescoping

GIVEN

$$A(n) := \sum_{k=1}^a \underbrace{\frac{S_1(k) + S_1(n) - S_1(k+n)}{kn(k+n+1)}}_{=: f(k)}.$$

FIND  $g(k)$  :

$$\boxed{g(k+1) - g(k)} = \boxed{f(k)}$$

for all  $0 \leq k \leq n$  and all  $n \geq 0$ .

## Telescoping

GIVEN

$$A(n) := \sum_{k=1}^a \underbrace{\frac{S_1(k) + S_1(n) - S_1(k+n)}{kn(k+n+1)}}_{=: f(k)}.$$

FIND  $g(k)$  :

$$\boxed{g(k+1) - g(k)} = \boxed{f(k)}$$

for all  $0 \leq k \leq n$  and all  $n \geq 0$ .

**no solution** 😞



## Zeilberger's creative telescoping paradigm

GIVEN

$$A(n) := \sum_{k=1}^a \underbrace{\frac{S_1(k) + S_1(n) - S_1(k+n)}{kn(k+n+1)}}_{=: f(n, k)}.$$

FIND  $g(n, k)$ 

$$\boxed{g(n, k+1) - g(n, k)} = \boxed{f(n, k)}$$

for all  $0 \leq k \leq n$  and all  $n \geq 0$ .**no solution** 

## Zeilberger's creative telescoping paradigm

GIVEN

$$A(n) := \sum_{k=1}^a \underbrace{\frac{S_1(k) + S_1(n) - S_1(k+n)}{kn(k+n+1)}}_{=: f(n, k)}.$$

FIND  $g(n, k)$  and  $c_0(n), c_1(n)$ :

$$\boxed{g(n, k+1) - g(n, k)} = \boxed{c_0(n)f(n, k) + c_1(n)f(n+1, k)}$$

for all  $0 \leq k \leq n$  and all  $n \geq 0$ .

## Zeilberger's creative telescoping paradigm

GIVEN

$$A(n) := \sum_{k=1}^a \underbrace{\frac{S_1(k) + S_1(n) - S_1(k+n)}{kn(k+n+1)}}_{=: f(n, k)}.$$

FIND  $g(n, k)$  and  $c_0(n), c_1(n)$ :

$$\boxed{g(n, k+1) - g(n, k)} = \boxed{c_0(n)f(n, k) + c_1(n)f(n+1, k)}$$

for all  $0 \leq k \leq n$  and all  $n \geq 0$ .

**Sigma computes:**  $c_0(n) = -n$ ,  $c_1(n) = (n+2)$  and

$$g(n, k) = \frac{kS_1(k) + (-n-1)S_1(n) - kS_1(k+n) - 2}{(k+n+1)(n+1)^2}$$

## Zeilberger's creative telescoping paradigm

GIVEN

$$A(n) := \sum_{k=1}^a \underbrace{\frac{S_1(k) + S_1(n) - S_1(k+n)}{kn(k+n+1)}}_{=: f(n, k)}.$$

FIND  $g(n, k)$  and  $c_0(n), c_1(n)$ :

$$\boxed{g(n, k+1) - g(n, k)} = \boxed{c_0(n)f(n, k) + c_1(n)f(n+1, k)}$$

for all  $0 \leq k \leq n$  and all  $n \geq 0$ .

Summing this equation over  $k$  from 1 to  $a$  gives:

$$\boxed{g(n, a+1) - g(n, 1)} = \boxed{\sum_{k=1}^a [c_0(n)f(n, k) + c_1(n)f(n+1, k)]}$$

## Zeilberger's creative telescoping paradigm

GIVEN

$$A(n) := \sum_{k=1}^a \underbrace{\frac{S_1(k) + S_1(n) - S_1(k+n)}{kn(k+n+1)}}_{=: f(n, k)}.$$

FIND  $g(n, k)$  and  $c_0(n), c_1(n)$ :

$$\boxed{g(n, k+1) - g(n, k)} = \boxed{c_0(n)f(n, k) + c_1(n)f(n+1, k)}$$

for all  $0 \leq k \leq n$  and all  $n \geq 0$ .

Summing this equation over  $k$  from 1 to  $a$  gives:

$$\boxed{g(n, a+1) - g(n, 1)} = \boxed{\sum_{k=1}^a c_0(n) f(n, k) + \sum_{k=1}^a c_1(n) f(n+1, k)}$$

## Zeilberger's creative telescoping paradigm

GIVEN

$$A(n) := \sum_{k=1}^a \underbrace{\frac{S_1(k) + S_1(n) - S_1(k+n)}{kn(k+n+1)}}_{=: f(n, k)}.$$

FIND  $g(n, k)$  and  $c_0(n), c_1(n)$ :

$$\boxed{g(n, k+1) - g(n, k)} = \boxed{c_0(n)f(n, k) + c_1(n)f(n+1, k)}$$

for all  $0 \leq k \leq n$  and all  $n \geq 0$ .

Summing this equation over  $k$  from 1 to  $a$  gives:

$$\boxed{g(n, a+1) - g(n, 1)} = \boxed{c_0(n) \sum_{k=1}^a f(n, k) + c_1(n) \sum_{k=1}^a f(n+1, k)}$$

## Zeilberger's creative telescoping paradigm

GIVEN

$$A(n) := \sum_{k=1}^a \underbrace{\frac{S_1(k) + S_1(n) - S_1(k+n)}{kn(k+n+1)}}_{=: f(n, k)}.$$

FIND  $g(n, k)$  and  $c_0(n), c_1(n)$ :

$$\boxed{g(n, k+1) - g(n, k)} = \boxed{c_0(n)f(n, k) + c_1(n)f(n+1, k)}$$

for all  $0 \leq k \leq n$  and all  $n \geq 0$ .

Summing this equation over  $k$  from 1 to  $a$  gives:

$$\boxed{g(n, a+1) - g(n, 1)} = \boxed{c_0(n)A(n) + c_1(n)A(n+1)}$$

## Zeilberger's creative telescoping paradigm

GIVEN

$$A(n) := \sum_{k=1}^a \underbrace{\frac{S_1(k) + S_1(n) - S_1(k+n)}{kn(k+n+1)}}_{=: f(n, k)}.$$

FIND  $g(n, k)$  and  $c_0(n), c_1(n)$ :

$$\boxed{g(n, k+1) - g(n, k)} = \boxed{c_0(n)f(n, k) + c_1(n)f(n+1, k)}$$

for all  $0 \leq k \leq n$  and all  $n \geq 0$ .

Summing this equation over  $k$  from 1 to  $a$  gives:

$$\begin{aligned} \boxed{g(n, a+1) - g(n, 1)} &= \boxed{c_0(n)A(n) + c_1(n)A(n+1)} \\ \parallel & \qquad \qquad \qquad \parallel \\ \frac{(a+1)(S_1(a)+S_1(n)-S_1(a+n))}{(n+1)^2(a+n+2)} & \qquad \qquad \qquad - nA(n) + (2+n)A(n+1) \\ + \frac{a(a+1)}{(n+1)^3(a+n+1)(a+n+2)} & \qquad \qquad \qquad \end{aligned}$$



$$(n+2)\mathbf{A}(n+1) - n\mathbf{A}(n) = \frac{(n+1)S_1(n) + 1}{(n+1)^3}$$

recurrence finder

$$A(n) = \sum_{k=1}^{\infty} \frac{S_1(k) + S_1(n) - S_1(k+n)}{kn(k+n+1)}$$

$$(n+2)\mathbf{A}(n+1) - n\mathbf{A}(n) = \frac{(n+1)S_1(n) + 1}{(n+1)^3}$$

recurrence solver

$$A(n) = \sum_{k=1}^{\infty} \frac{S_1(k) + S_1(n) - S_1(k+n)}{kn(k+n+1)}$$

$\in$

$$\left\{ c \times \frac{1}{n(n+1)} + \frac{S_1(n)^2 + S_2(n)}{2n(n+1)} \mid c \in \mathbb{R} \right\}$$

where

$$S_1(n) = \sum_{i=1}^n \frac{1}{i}$$

$$S_2(n) = \sum_{i=1}^n \frac{1}{i^2}$$

$$(n+2)\mathbf{A}(n+1) - n\mathbf{A}(n) = \frac{(n+1)S_1(n) + 1}{(n+1)^3}$$

## Summation package Sigma

(based on difference field/ring algorithms/theory

see, e.g., Abramov, Karr 1981, Bronstein 2000, Schneider 2001/2004/2005a-c/2007/2008/2010a-c)

$$A(n) = \sum_{k=1}^{\infty} \frac{S_1(k) + S_1(n) - S_1(k+n)}{kn(k+n+1)}$$

$$= 0 \times \frac{1}{n(n+1)} + \frac{S_1(n)^2 + S_2(n)}{2n(n+1)}$$

where

$$S_1(n) = \sum_{i=1}^n \frac{1}{i}$$

$$S_2(n) = \sum_{i=1}^n \frac{1}{i^2}$$

$$\text{ln[12]:= mySum} = \sum_{k=1}^a \frac{S[1, k] + S[1, n] - S[1, k + n]}{kn(k + n + 1)};$$

$$\text{In[12]:= mySum} = \sum_{k=1}^a \frac{S[1, k] + S[1, n] - S[1, k + n]}{kn(k + n + 1)};$$

## Compute a recurrence

$$\text{In[13]:= rec} = \text{GenerateRecurrence}[\text{mySum}, n][[1]]$$

$$\text{Out[13]= } n\text{SUM}[n] + (1+n)(2+n)\text{SUM}[n+1] == \frac{(a+1)(S[1,a]+S[1,n]-S[1,a+n])}{(n+1)^2(a+n+2)n!} + \frac{a(a+1)}{(n+1)^3(a+n+1)(a+n+2)n!}$$

$$\text{In[12]:= mySum} = \sum_{k=1}^a \frac{S[1, k] + S[1, n] - S[1, k + n]}{kn(k + n + 1)};$$

## Compute a recurrence

$$\text{In[13]:= rec} = \text{GenerateRecurrence}[\text{mySum}, n][[1]]$$

$$\text{Out[13]= } n\text{SUM}[n] + (1+n)(2+n)\text{SUM}[n+1] == \frac{(a+1)(S[1,a]+S[1,n]-S[1,a+n])}{(n+1)^2(a+n+2)n!} + \frac{a(a+1)}{(n+1)^3(a+n+1)(a+n+2)n!}$$

$$\text{In[14]:= rec} = \text{LimitRec}[\text{rec}, \text{SUM}[n], \{n\}, a]$$

$$\text{Out[14]= } -n\text{SUM}[n] + (1+n)(2+n)\text{SUM}[n+1] == \frac{(n+1)S[1,n] + 1}{(n+1)^3}$$

$$\text{In[12]:= mySum} = \sum_{k=1}^a \frac{S[1, k] + S[1, n] - S[1, k + n]}{kn(k + n + 1)};$$

## Compute a recurrence

$$\text{In[13]:= rec} = \text{GenerateRecurrence}[\text{mySum}, n][[1]]$$

$$\text{Out[13]= } n\text{SUM}[n] + (1+n)(2+n)\text{SUM}[n+1] == \frac{(a+1)(S[1,a]+S[1,n]-S[1,a+n])}{(n+1)^2(a+n+2)n!} + \frac{a(a+1)}{(n+1)^3(a+n+1)(a+n+2)n!}$$

$$\text{In[14]:= rec} = \text{LimitRec}[\text{rec}, \text{SUM}[n], \{n\}, a]$$

$$\text{Out[14]= } -n\text{SUM}[n] + (1+n)(2+n)\text{SUM}[n+1] == \frac{(n+1)S[1,n] + 1}{(n+1)^3}$$

## Solve a recurrence

$$\text{In[15]:= recSol} = \text{SolveRecurrence}[\text{rec}, \text{SUM}[n]]$$

$$\text{Out[15]= } \left\{ \left\{ 0, \frac{1}{n(n+1)} \right\}, \left\{ 1, \frac{S[1,n]^2 + \sum_{i=1}^n \frac{1}{i^2}}{2n(n+1)} \right\} \right\}$$

$$\text{In[12]:= mySum} = \sum_{k=1}^a \frac{S[1, k] + S[1, n] - S[1, k + n]}{kn(k + n + 1)};$$

## Compute a recurrence

$$\text{In[13]:= rec} = \text{GenerateRecurrence}[\text{mySum}, n][[1]]$$

$$\text{Out[13]= } n\text{SUM}[n] + (1+n)(2+n)\text{SUM}[n+1] == \frac{(a+1)(S[1,a]+S[1,n]-S[1,a+n])}{(n+1)^2(a+n+2)n!} + \frac{a(a+1)}{(n+1)^3(a+n+1)(a+n+2)n!}$$

$$\text{In[14]:= rec} = \text{LimitRec}[\text{rec}, \text{SUM}[n], \{n\}, a]$$

$$\text{Out[14]= } -n\text{SUM}[n] + (1+n)(2+n)\text{SUM}[n+1] == \frac{(n+1)S[1,n] + 1}{(n+1)^3}$$

## Solve a recurrence

$$\text{In[15]:= recSol} = \text{SolveRecurrence}[\text{rec}, \text{SUM}[n]]$$

$$\text{Out[15]= } \left\{ \left\{ 0, \frac{1}{n(n+1)} \right\}, \left\{ 1, \frac{S[1,n]^2 + \sum_{i=1}^n \frac{1}{i^2}}{2n(n+1)} \right\} \right\}$$

## Combine the solutions

$$\text{In[16]:= FindLinearCombination}[\text{recSol}, \{1, \{1/2\}, n, 2\}$$

$$\text{Out[16]= } \frac{S[1,n]^2 + \sum_{i=1}^n \frac{1}{i^2}}{2n(n+1)}$$



## A warm-up example: simplify

$$\sum_{k=0}^{\infty} \sum_{j=0}^{\infty} \left( \frac{(2j+k+n+2)j!k!(j+k+n)!}{(j+k+1)(j+n+1)(j+k+1)!(j+n+1)!(k+n+1)!} \right. \\ \left. + \frac{j!k!(j+k+n)!(-S_1(j) + S_1(j+k) + S_1(j+n) - S_1(j+k+n))}{(j+k+1)!(j+n+1)!(k+n+1)!} \right) \\ \underbrace{\hspace{15em}}_{f(j)}$$

$$\sum_{k=1}^{\infty} \sum_{j=0}^{\infty} f(j) = \frac{1}{n!} \sum_{k=1}^{\infty} \frac{S_1(k) + S_1(n) - S_1(k+n)}{kn(k+n+1)} \\ = \frac{1}{n!} \frac{S_1(n)^2 + S_2(n)}{2n(n+1)}$$

where

$$S_1(n) = \sum_{i=1}^n \frac{1}{i} \qquad S_2(n) = \sum_{i=1}^n \frac{1}{i^2}$$

## A warm-up example: simplify

$$\sum_{k=0}^{\infty} \sum_{j=0}^{\infty} \left( \frac{(2j+k+n+2)j!k!(j+k+n)!}{(j+k+1)(j+n+1)(j+k+1)!(j+n+1)!(k+n+1)!} \right. \\ \left. + \frac{j!k!(j+k+n)!(-S_1(j) + S_1(j+k) + S_1(j+n) - S_1(j+k+n))}{(j+k+1)!(j+n+1)!(k+n+1)!} \right) \\ \underbrace{\hspace{15em}}_{f(n, k, j)}$$

$$\sum_{k=0}^{\infty} \sum_{j=0}^{\infty} f(n, k, j) = \frac{S_1(n)^2 + 3S_2(n)}{2n(n+1)!}$$

where

$$S_1(n) = \sum_{i=1}^n \frac{1}{i} \qquad S_2(n) = \sum_{i=1}^n \frac{1}{i^2}$$

# 1. Creative telescoping (for the special case of hypergeometric terms see Zeilberger's algorithm (1991))

GIVEN a **definite** sum

$$A(n) = \sum_{k=0}^n f(n, k);$$

$f(n, k)$ : indefinite nested product-sum in  $k$ ;  
 $n$ : extra parameter

FIND a **recurrence** for  $A(n)$

## 1. Creative telescoping (for the special case of hypergeometric terms see Zeilberger's algorithm (1991))

GIVEN a definite sum

$$A(n) = \sum_{k=0}^n f(n, k);$$

$f(n, k)$ : indefinite nested product-sum in  $k$ ;  
 $n$ : extra parameter

FIND a recurrence for  $A(n)$ 

## 2. Recurrence solving

GIVEN a recurrence

$a_0(n), \dots, a_d(n), h(n)$ :  
 indefinite nested product-sum expressions.

$$a_0(n)A(n) + \dots + a_d(n)A(n+d) = h(n);$$

FIND all solutions expressible by indefinite nested products/sums

(Abramov/Bronstein/Petkovšek/CS, in preparation)

## 1. Creative telescoping (for the special case of hypergeometric terms see Zeilberger's algorithm (1991))

GIVEN a definite sum

$$A(n) = \sum_{k=0}^n f(n, k);$$

$f(n, k)$ : indefinite nested product-sum in  $k$ ;  
 $n$ : extra parameter

FIND a recurrence for  $A(n)$ 

## 2. Recurrence solving

GIVEN a recurrence

$a_0(n), \dots, a_d(n), h(n)$ :  
 indefinite nested product-sum expressions.

$$a_0(n)A(n) + \dots + a_d(n)A(n+d) = h(n);$$

FIND all solutions expressible by indefinite nested products/sums

(Abramov/Bronstein/Petkovšek/CS, in preparation)

## 3. Find a "closed form"

$A(n)$ =combined solutions in terms of indefinite nested sums.

$$\sum_{j=0}^{n-2} \sum_{r=0}^{j+1} \sum_{s=0}^{n-j+r-2} \frac{(-1)^{r+s} \binom{j+1}{r} \binom{-j+n+r-2}{s} (-j+n-2)! r!}{(n-s)(s+1)(-j+n+r)!}$$

Simple sum

$$\sum_{j=0}^{n-2} \sum_{r=0}^{j+1} \sum_{s=0}^{n-j+r-2} \frac{(-1)^{r+s} \binom{j+1}{r} \binom{-j+n+r-2}{s} (-j+n-2)! r!}{(n-s)(s+1)(-j+n+r)!}$$

||

$$\sum_{j=0}^{n-2} \sum_{r=0}^{j+1} \boxed{\sum_{s=0}^{n-j+r-2} \frac{(-1)^{r+s} \binom{j+1}{r} \binom{-j+n+r-2}{s} (-j+n-2)! r!}{(n-s)(s+1)(-j+n+r)!}}$$

$$\sum_{j=0}^{n-2} \sum_{r=0}^{j+1} \sum_{s=0}^{n-j+r-2} \frac{(-1)^{r+s} \binom{j+1}{r} \binom{-j+n+r-2}{s} (-j+n-2)! r!}{(n-s)(s+1)(-j+n+r)!}$$

||

$$\sum_{j=0}^{n-2} \sum_{r=0}^{j+1} \boxed{\sum_{s=0}^{n-j+r-2} \frac{(-1)^{r+s} \binom{j+1}{r} \binom{-j+n+r-2}{s} (-j+n-2)! r!}{(n-s)(s+1)(-j+n+r)!}}$$

||

$$\boxed{\binom{j+1}{r} \left( \frac{(-1)^r (-j+n-2)! r!}{(n+1)(-j+n+r-1)(-j+n+r)!} + \frac{(-1)^{n+r} (j+1)! (-j+n-2)! (-j+n-1)_r r!}{(n-1)n(n+1)(-j+n+r)! (-j-1)_r (2-n)_j} \right)}$$



$$\sum_{j=0}^{n-2} \sum_{r=0}^{j+1} \sum_{s=0}^{n-j+r-2} \frac{(-1)^{r+s} \binom{j+1}{r} \binom{-j+n+r-2}{s} (-j+n-2)! r!}{(n-s)(s+1)(-j+n+r)!}$$

||

$$\sum_{j=0}^{n-2} \left( \sum_{r=0}^{j+1} \binom{j+1}{r} \left( \frac{(-1)^r (-j+n-2)! r!}{(n+1)(-j+n+r-1)(-j+n+r)!} + \frac{(-1)^{n+r} (j+1)! (-j+n-2)! (-j+n-1)_r r!}{(n-1)n(n+1)(-j+n+r)! (-j-1)_r (2-n)_j} \right) \right)$$

$$\sum_{j=0}^{n-2} \sum_{r=0}^{j+1} \sum_{s=0}^{n-j+r-2} \frac{(-1)^{r+s} \binom{j+1}{r} \binom{-j+n+r-2}{s} (-j+n-2)! r!}{(n-s)(s+1)(-j+n+r)!}$$

$$\parallel$$

$$\sum_{j=0}^{n-2} \left( \sum_{r=0}^{j+1} \binom{j+1}{r} \left( \frac{(-1)^r (-j+n-2)! r!}{(n+1)(-j+n+r-1)(-j+n+r)!} + \frac{(-1)^{n+r} (j+1)! (-j+n-2)! (-j+n-1)_r r!}{(n-1)n(n+1)(-j+n+r)! (-j-1)_r (2-n)_j} \right) \right)$$

$$\parallel$$

$$\left( \frac{n^2 - n + 1}{(n-1)^2 n^2 (n+1)(2-n)_j} + \frac{\sum_{i=1}^j \frac{(2-n)_i}{(-i+n-1)^2 (i+1)!}}{(n+1)(2-n)_j} + \frac{(-1)^{j+n} (-j-2)(-j+n-2)!}{(j-n+1)(n+1)^2 n!} \right) (j+1)! - \frac{1}{(n+1)^2 (-j+n-1)}$$

$$\sum_{j=0}^{n-2} \sum_{r=0}^{j+1} \sum_{s=0}^{n-j+r-2} \frac{(-1)^{r+s} \binom{j+1}{r} \binom{-j+n+r-2}{s} (-j+n-2)! r!}{(n-s)(s+1)(-j+n+r)!}$$

||

$$\sum_{j=0}^{n-2} \left( \left( \frac{n^2 - n + 1}{(n-1)^2 n^2 (n+1)(2-n)_j} + \frac{\sum_{i=1}^j \frac{(2-n)_i}{(-i+n-1)^2 (i+1)!}}{(n+1)(2-n)_j} + \frac{(-1)^{j+n} (-j-2)(-j+n-2)!}{(j-n+1)(n+1)^2 n!} \right) (j+1)! - \frac{1}{(n+1)^2 (-j+n-1)} \right)$$

$$\sum_{j=0}^{n-2} \sum_{r=0}^{j+1} \sum_{s=0}^{n-j+r-2} \frac{(-1)^{r+s} \binom{j+1}{r} \binom{-j+n+r-2}{s} (-j+n-2)! r!}{(n-s)(s+1)(-j+n+r)!}$$

||

$$\sum_{j=0}^{n-2} \left( \left( \frac{n^2 - n + 1}{(n-1)^2 n^2 (n+1)(2-n)_j} + \frac{\sum_{i=1}^j \frac{(2-n)_i}{(-i+n-1)^2 (i+1)!}}{(n+1)(2-n)_j} + \frac{(-1)^{j+n} (-j-2)(-j+n-2)!}{(j-n+1)(n+1)^2 n!} \right) (j+1)! - \frac{1}{(n+1)^2 (-j+n-1)} \right)$$

||

$$\frac{-n^2 - n - 1}{n^2(n+1)^3} + \frac{(-1)^n (n^2 + n + 1)}{n^2(n+1)^3} - \frac{2S_{-2}(n)}{n+1} + \frac{S_1(n)}{(n+1)^2} + \frac{S_2(n)}{-n-1}$$

Note:  $S_a(n) = \sum_{i=1}^N \frac{\text{sign}(a)^i}{i^{|a|}}$ ,  $a \in \mathbb{Z} \setminus \{0\}$ .

In[1]:= << **Sigma.m**

Sigma - A summation package by Carsten Schneider © RISC-Linz

In[2]:= << **HarmonicSums.m**

HarmonicSums by Jakob Ablinger © RISC-Linz

In[3]:= << **EvaluateMultiSums.m**

EvaluateMultiSums by Carsten Schneider © RISC-Linz

In[1]:= << **Sigma.m**

Sigma - A summation package by Carsten Schneider © RISC-Linz

In[2]:= << **HarmonicSums.m**

HarmonicSums by Jakob Ablinger © RISC-Linz

In[3]:= << **EvaluateMultiSums.m**

EvaluateMultiSums by Carsten Schneider © RISC-Linz

$$\text{In[4]:= mySum} = \sum_{j=0}^{n-2} \sum_{r=0}^{j+1} \sum_{s=0}^{n-j+r-2} \frac{(-1)^{r+s} \binom{j+1}{r} \binom{-j+n+r-2}{s} (-j+n-2)! r!}{(n-s)(s+1)(-j+n+r)!};$$

In[5]:= **EvaluateMultiSum**[mySum, {}, {n}, {1}]

In[1]:= << **Sigma.m**

Sigma - A summation package by Carsten Schneider © RISC-Linz

In[2]:= << **HarmonicSums.m**

HarmonicSums by Jakob Ablinger © RISC-Linz

In[3]:= << **EvaluateMultiSums.m**

EvaluateMultiSums by Carsten Schneider © RISC-Linz

$$\text{In[4]:= mySum} = \sum_{j=0}^{n-2} \sum_{r=0}^{j+1} \sum_{s=0}^{n-j+r-2} \frac{(-1)^{r+s} \binom{j+1}{r} \binom{-j+n+r-2}{s} (-j+n-2)! r!}{(n-s)(s+1)(-j+n+r)!};$$


In[5]:= **EvaluateMultiSum**[mySum, {}, {n}, {1}]

$$\text{Out[5]=} \frac{-n^2 - n - 1}{n^2(n+1)^3} + \frac{(-1)^n (n^2 + n + 1)}{n^2(n+1)^3} - \frac{2S[-2, n]}{n+1} + \frac{S[1, n]}{(n+1)^2} + \frac{S[2, n]}{-n-1}$$

Application: The simplification of  
Feynman integrals




## Evaluation of Feynman Integrals



Behavior of particles

## Evaluation of Feynman Integrals



Behavior of particles




$$\int \Phi(N, \epsilon, x) dx$$


Feynman integrals

$$\int_0^1 x dx = ?$$

$$\int_0^1 x dx =$$




$$\int_0^1 x^1 dx =$$




$$\int_0^1 x^2 dx = ?$$


$$\int_0^1 x^1 dx =$$




$$\int_0^1 x^2 dx =$$



$$\int_0^1 x^1 dx =$$




$$\int_0^1 x^2 dx =$$




$$\int_0^1 x^3 dx = ?$$


$$\int_0^1 x^1 dx =$$



$$\int_0^1 x^2 dx =$$




$$\int_0^1 x^3 dx =$$







$$\int_0^1 x^1 dx =$$



$$\int_0^1 x^2 dx =$$



$$\int_0^1 x^3 dx =$$



$$\int_0^1 x^N dx = \frac{1}{N+1}$$

für  $N = 1, 2, 3, 4, \dots$

## Feynman integrals

$$\int_0^1 x^N dx$$

## Feynman integrals

$$\int_0^1 x^N (1+x)^N dx$$

## Feynman integrals

$$\int_0^1 \frac{x^N (1+x)^N}{(1-x)^{1+\varepsilon}} dx$$

## Feynman integrals

$$\int_0^1 \int_0^1 \frac{x_1^N (1+x_1)^N}{(1-x_1)^{1+\varepsilon}} \dots dx_1 dx_2$$

## Feynman integrals

$$\int_0^1 \int_0^1 \int_0^1 \frac{x_1^N (1+x_1)^N}{(1-x_1)^{1+\varepsilon}} \dots dx_1 dx_2 dx_3$$

## Feynman integrals

$$\int_0^1 \int_0^1 \int_0^1 \int_0^1 \frac{x_1^N (1+x_1)^N}{(1-x_1)^{1+\varepsilon}} \dots dx_1 dx_2 dx_3 dx_4$$

## Feynman integrals

$$\int_0^1 \int_0^1 \int_0^1 \int_0^1 \int_0^1 \frac{x_1^N (1+x_1)^N}{(1-x_1)^{1+\varepsilon}} \dots dx_1 dx_2 dx_3 dx_4 dx_5$$




## Feynman integrals

$$\int_0^1 \int_0^1 \int_0^1 \int_0^1 \int_0^1 \int_0^1 \frac{x_1^N (1+x_1)^N}{(1-x_1)^{1+\varepsilon}} \dots dx_1 dx_2 dx_3 dx_4 dx_5 dx_6$$

## Feynman integrals

$$\sum_{j=0}^{N-3} \sum_{k=0}^j \binom{N-1}{j+2} \binom{j+1}{k+1} \\ \times \int_0^1 \int_0^1 \int_0^1 \int_0^1 \int_0^1 \int_0^1 \frac{x_1^N (1+x_1)^{N-j+k}}{(1-x_1)^{1+\varepsilon}} \dots dx_1 dx_2 dx_3 dx_4 dx_5 dx_6$$


## Feynman integrals



a 3-loop massive ladder diagram [arXiv:1509.08324]

$$\begin{aligned}
 & \sum_{j=0}^{N-3} \sum_{k=0}^j \binom{N-1}{j+2} \binom{j+1}{k+1} \quad || \\
 & \times \int_0^1 \int_0^1 \int_0^1 \int_0^1 \int_0^1 \int_0^1 \theta(1-x_5-x_6)(1-x_2)(1-x_4)x_2^{-\varepsilon} \\
 & (1-x_2)^{-\varepsilon} x_4^{\varepsilon/2-1} (1-x_4)^{\varepsilon/2-1} x_5^{\varepsilon-1} x_6^{-\varepsilon p/2} \\
 & \left[ \begin{aligned}
 & [-x_3(1-x_4) - x_4(1-x_5-x_6 + x_5x_1 + x_6x_3)]^k \\
 & + [x_3(1-x_4) - (1-x_4)(1-x_5-x_6 + x_5x_1 + x_6x_3)]^k
 \end{aligned} \right] \\
 & \times (1-x_5-x_6 + x_5x_1 + x_6x_3)^{j-k} (1-x_2)^{N-3-j} \\
 & \times [x_1 - (1-x_5-x_6) - x_5x_1 - x_6x_3]^{N-3-j} dx_1 dx_2 dx_3 dx_4 dx_5 dx_6
 \end{aligned}$$

## Evaluation of Feynman Integrals




Behavior of particles



$$\int \Phi(N, \epsilon, x) dx$$

Feynman integrals

# Evaluation of Feynman Integrals



Behavior of particles



$$\int \Phi(N, \epsilon, x) dx$$


Feynman integrals

**DESY**  
(J. Blümlein)

$$\sum f(N, \epsilon, k)$$

complicated  
multi-sums

# Evaluation of Feynman Integrals



Behavior of particles



$$\int \Phi(N, \epsilon, x) dx$$

Feynman integrals

**DESY**  
(J. Blümlein)



$$\sum f(N, \epsilon, k)$$

complicated  
multi-sums

expression in  
special functions




**RISC**  
(Sigma-package)

Example 1:

massive 3-loop ladder integrals


## Feynman integrals




a 3-loop massive ladder diagram [arXiv:1509.08324]

$$\begin{aligned}
 & \sum_{j=0}^{N-3} \sum_{k=0}^j \binom{N-1}{j+2} \binom{j+1}{k+1} \quad || \\
 & \times \int_0^1 \int_0^1 \int_0^1 \int_0^1 \int_0^1 \int_0^1 \theta(1-x_5-x_6)(1-x_2)(1-x_4)x_2^{-\varepsilon} \\
 & (1-x_2)^{-\varepsilon} x_4^{\varepsilon/2-1} (1-x_4)^{\varepsilon/2-1} x_5^{\varepsilon-1} x_6^{-\varepsilon p/2} \\
 & \left[ \begin{aligned}
 & [-x_3(1-x_4) - x_4(1-x_5-x_6 + x_5x_1 + x_6x_3)]^k \\
 & + [x_3(1-x_4) - (1-x_4)(1-x_5-x_6 + x_5x_1 + x_6x_3)]^k
 \end{aligned} \right] \\
 & \times (1-x_5-x_6 + x_5x_1 + x_6x_3)^{j-k} (1-x_2)^{N-3-j} \\
 & \times [x_1 - (1-x_5-x_6) - x_5x_1 - x_6x_3]^{N-3-j} dx_1 dx_2 dx_3 dx_4 dx_5 dx_6
 \end{aligned}$$





$$= F_{-3}(N)\varepsilon^{-3} + F_{-2}(N)\varepsilon^{-2} + F_{-1}(N)\varepsilon^{-1} + \boxed{F_0(N)}$$



$$= F_{-3}(N)\varepsilon^{-3} + F_{-2}(N)\varepsilon^{-2} + F_{-1}(N)\varepsilon^{-1} + \boxed{F_0(N)}$$

Simplify

||

$$\sum_{j=0}^{N-3} \sum_{k=0}^j \sum_{l=0}^k \sum_{q=0}^{-j+N-3} \sum_{s=1}^{-l+N-q-3} \sum_{r=0}^{-l+N-q-s-3} (-1)^{-j+k-l+N-q-3} \times$$

$$\times \frac{(j+1)(k)(N-1)(-j+N-3)(-l+N-q-3)(-l+N-q-s-3)r!(-l+N-q-r-s-3)!(s-1)!}{(-l+N-q-2)!(-j+N-1)(N-q-r-s-2)(q+s+1)}$$

$$\left[ 4S_1(-j+N-1) - 4S_1(-j+N-2) - 2S_1(k) \right.$$

$$\left. - (S_1(-l+N-q-2) + S_1(-l+N-q-r-s-3) - 2S_1(r+s)) \right.$$

$$\left. + 2S_1(s-1) - 2S_1(r+s) \right] + \mathbf{3 \text{ further 6-fold sums}}$$

$$\boxed{F_0(N)} =$$

$$\begin{aligned} & \frac{7}{12}S_1(N)^4 + \frac{(17N+5)S_1(N)^3}{3N(N+1)} + \left( \frac{35N^2-2N-5}{2N^2(N+1)^2} + \frac{13S_2(N)}{2} + \frac{5(-1)^N}{2N^2} \right) S_1(N)^2 \\ & + \left( -\frac{4(13N+5)}{N^2(N+1)^2} + \left( \frac{4(-1)^N(2N+1)}{N(N+1)} - \frac{13}{N} \right) S_2(N) + \left( \frac{29}{3} - (-1)^N \right) S_3(N) \right. \\ & + \left( 2 + 2(-1)^N \right) S_{2,1}(N) - 28S_{-2,1}(N) + \frac{20(-1)^N}{N^2(N+1)} S_1(N) + \left( \frac{3}{4} + (-1)^N \right) S_2(N)^2 \\ & - 2(-1)^N S_{-2}(N)^2 + S_{-3}(N) \left( \frac{2(3N-5)}{N(N+1)} + (26 + 4(-1)^N) S_1(N) + \frac{4(-1)^N}{N+1} \right) \\ & + \left( \frac{(-1)^N(5-3N)}{2N^2(N+1)} - \frac{5}{2N^2} \right) S_2(N) + S_{-2}(N) (10S_1(N)^2 + \left( \frac{8(-1)^N(2N+1)}{N(N+1)} \right. \\ & + \left. \frac{4(3N-1)}{N(N+1)} \right) S_1(N) + \frac{8(-1)^N(3N+1)}{N(N+1)^2} + (-22 + 6(-1)^N) S_2(N) - \frac{16}{N(N+1)} \\ & + \left( \frac{(-1)^N(9N+5)}{N(N+1)} - \frac{29}{3N} \right) S_3(N) + \left( \frac{19}{2} - 2(-1)^N \right) S_4(N) + (-6 + 5(-1)^N) S_{-4}(N) \\ & + \left( -\frac{2(-1)^N(9N+5)}{N(N+1)} - \frac{2}{N} \right) S_{2,1}(N) + (20 + 2(-1)^N) S_{2,-2}(N) + (-17 + 13(-1)^N) S_{3,1}(N) \\ & - \frac{8(-1)^N(2N+1) + 4(9N+1)}{N(N+1)} S_{-2,1}(N) - (24 + 4(-1)^N) S_{-3,1}(N) + (3 - 5(-1)^N) S_{2,1,1}(N) \\ & + 32S_{-2,1,1}(N) + \left( \frac{3}{2}S_1(N)^2 - \frac{3S_1(N)}{N} + \frac{3}{2}(-1)^N S_{-2}(N) \right) \zeta(2) \end{aligned}$$

$$\boxed{F_0(N)} =$$

$$\begin{aligned} & \frac{7}{12} S_1(N) + \frac{(17N+5)S_1(N)^3}{2N^2(N+1)^2} + \left( \frac{35N^2 - 2N - 5}{2N^2(N+1)^2} + \frac{13S_2(N)}{2} + \frac{5(-1)^N}{2N^2} \right) S_1(N)^2 \\ & + \left( - \frac{S_1(N)}{N(N+1)} + \frac{1}{i} \right) S_1(N) = \sum_{i=1}^N \frac{1}{i} \left( \frac{1}{N(N+1)} - \frac{13}{N} \right) S_2(N) + \left( \frac{29}{3} - (-1)^N \right) S_3(N) \\ & + \left( 2 + 2(-1)^N \right) S_{2,1}(N) - 28S_{-2,1}(N) + \frac{20(-1)^N}{N^2(N+1)} S_1(N) + \left( \frac{3}{4} + (-1)^N \right) S_2(N)^2 \\ & - 2(-1)^N S_{-2}(N)^2 + S_{-3}(N) \left( \frac{2(3N-5)}{N(N+1)} + (26 + 4(-1)^N) S_1(N) + \frac{4(-1)^N}{N+1} \right) \\ & + \left( \frac{(-1)^N(5-3N)}{2N^2(N+1)} - \frac{5}{2N^2} \right) S_2(N) + S_{-2}(N) (10S_1(N)^2 + \left( \frac{8(-1)^N(2N+1)}{N(N+1)} \right. \\ & \left. + \frac{4(3N-1)}{N(N+1)} \right) S_1(N) + \frac{8(-1)^N(3N+1)}{N(N+1)^2} + (-22 + 6(-1)^N) S_2(N) - \frac{16}{N(N+1)} \\ & + \left( \frac{(-1)^N(9N+5)}{N(N+1)} - \frac{29}{3N} \right) S_3(N) + \left( \frac{19}{2} - 2(-1)^N \right) S_4(N) + (-6 + 5(-1)^N) S_{-4}(N) \\ & + \left( - \frac{2(-1)^N(9N+5)}{N(N+1)} - \frac{2}{N} \right) S_{2,1}(N) + (20 + 2(-1)^N) S_{2,-2}(N) + (-17 + 13(-1)^N) S_{3,1}(N) \\ & - \frac{8(-1)^N(2N+1) + 4(9N+1)}{N(N+1)} S_{-2,1}(N) - (24 + 4(-1)^N) S_{-3,1}(N) + (3 - 5(-1)^N) S_{2,1,1}(N) \\ & + 32S_{-2,1,1}(N) + \left( \frac{3}{2} S_1(N)^2 - \frac{3S_1(N)}{N} + \frac{3}{2} (-1)^N S_{-2}(N) \right) \zeta(2) \end{aligned}$$

$$F_0(N) =$$

$$\begin{aligned}
& \frac{7}{12} S_1(N) + \frac{(17N+5)S_1(N)^3}{2N^2(N+1)^2} + \left( \frac{35N^2 - 2N - 5}{2N^2(N+1)^2} + \frac{13S_2(N)}{2} + \frac{5(-1)^N}{2N^2} \right) S_1(N)^2 \\
& + \left( -\frac{1}{N(N+1)} \right) S_1(N) = \sum_{i=1}^N \frac{1}{i} (-1)^N (2N+1) - \frac{13}{N} S_2(N) + \left( \frac{29}{2} - (-1)^N \right) S_3(N) \\
& + (2 + 2(-1)^N) S_{2,1}(N) - 28S_{-2,1}(N) + \frac{20(-1)^N}{N^2(N+1)} S_2(N) = \sum_{i=1}^N \frac{1}{i^2} (-1)^N S_2(N)^2 \\
& - 2(-1)^N S_{-2}(N)^2 + S_{-3}(N) \left( \frac{2(3N-5)}{N(N+1)} + (26 + 4(-1)^N) \frac{1}{N+1} \right) \\
& + \left( \frac{(-1)^N(5-3N)}{2N^2(N+1)} - \frac{5}{2N^2} \right) S_2(N) + S_{-2}(N) (10S_1(N)^2 + \frac{8(-1)^N(2N+1)}{N(N+1)}) \\
& + \frac{4(3N-1)}{N(N+1)} S_1(N) + \frac{8(-1)^N(3N+1)}{N(N+1)^2} + (-22 + 6(-1)^N) S_2(N) - \frac{16}{N(N+1)} \\
& + \left( \frac{(-1)^N(9N+5)}{N(N+1)} - \frac{29}{3N} \right) S_3(N) + \left( \frac{19}{2} - 2(-1)^N \right) S_4(N) + (-6 + 5(-1)^N) S_{-4}(N) \\
& + \left( -\frac{2(-1)^N(9N+5)}{N(N+1)} - \frac{2}{N} \right) S_{2,1}(N) + (20 + 2(-1)^N) S_{2,-2}(N) + (-17 + 13(-1)^N) S_{3,1}(N) \\
& - \frac{8(-1)^N(2N+1) + 4(9N+1)}{N(N+1)} S_{-2,1}(N) - (24 + 4(-1)^N) S_{-3,1}(N) + (3 - 5(-1)^N) S_{2,1,1}(N) \\
& + 32S_{-2,1,1}(N) + \left( \frac{3}{2} S_1(N)^2 - \frac{3S_1(N)}{N} + \frac{3}{2} (-1)^N S_{-2}(N) \right) \zeta(2)
\end{aligned}$$


$$\boxed{F_0(N)} =$$

$$\begin{aligned}
 & \frac{7}{12} S_1(N)^3 + \left( \frac{(17N+5)S_1(N)^3}{2N^2(N+1)^2} + \frac{35N^2 - 2N - 5}{2} + \frac{13S_2(N)}{2} + \frac{5(-1)^N}{2N^2} \right) S_1(N)^2 \\
 & + \left( -\frac{1}{N(N+1)} \right) S_1(N) = \sum_{i=1}^N \frac{1}{i} (-1)^N (2N+1) - \frac{13}{N} S_2(N) + \left( \frac{29}{2} - (-1)^N \right) S_3(N) \\
 & + (2 + 2(-1)^N) S_{2,1}(N) - 28S_{-2,1}(N) + \frac{20(-1)^N}{N^2(N+1)} S_2(N) = \sum_{i=1}^N \frac{1}{i^2} (-1)^N S_2(N)^2 \\
 & - 2(-1)^N S_{-2}(N)^2 + S_{-3}(N) \left( \frac{2(3N-5)}{N(N+1)} + (26 + 4(-1)^N) \frac{(-1)^N}{N+1} \right) \\
 & + \left( \frac{(-1)^N}{2N^2} \right)^2 + \left( \frac{8(-1)^N(2N+1)}{N(N+1)} \right) \\
 & + \frac{4(3N-5)}{N(N+1)} (-1)^N S_2(N) - \frac{16}{N(N+1)} \\
 & + \left( \frac{(-1)^N}{N} \right) S_{-2,1,1}(N) = \sum_{i=1}^N \frac{(-1)^i \sum_{j=1}^i \frac{1}{k}}{i^2} S_{-2,1,1}(N) + (-6 + 5(-1)^N) S_{-4}(N) \\
 & + \left( -\frac{2(-1)^N}{N(N+1)} \right) S_{2,-2}(N) + (-17 + 13(-1)^N) S_{3,1}(N) \\
 & - \frac{8(-1)^N(2N+1) + 4(9N+1)}{N(N+1)} S_{-2,1}(N) - (24 + 4(-1)^N) S_{-3,1}(N) + (3 - 5(-1)^N) S_{2,1,1}(N) \\
 & + 32S_{-2,1,1}(N) + \left( \frac{3}{2} S_1(N)^2 - \frac{3S_1(N)}{N} + \frac{3}{2} (-1)^N S_{-2}(N) \right) \zeta(2)
 \end{aligned}$$

Example 2:

2-mass 3-loop Feynman integrals

Example: a 2-mass 3-loop Feynman integral [arXiv:1804.02226]  
(arose in the calculation of the gluonic operator matrix element  $A_{gg,Q}^{(3)}$ )




All diagrams are produced with axodraw (J. Vermaseren).



# Example: a 2-mass 3-loop Feynman integral [arXiv:1804.02226]

(arose in the calculation of the gluonic operator matrix element  $A_{gg,Q}^{(3)}$ )



Mellin-Barnes-  
and  ${}_pF_q$ -technologies


→

expression (95 MB) with

- 150 single sums
- 1000 double sums
- 12160 triple sums
- 1555 quadruple sums

# Example: a 2-mass 3-loop Feynman integral [arXiv:1804.02226]

(arose in the calculation of the gluonic operator matrix element  $A_{gg,Q}^{(3)}$ )



Mellin-Barnes-  
and  $pF_q$ -technologies

expression (95 MB) with


- 150 single sums
- 1000 double sums
- 12160 triple sums
- 1555 quadruple sums

Typical triple sum:

$$\sum_{j=0}^N \sum_{i=0}^j \sum_{k=0}^i \frac{(4+\varepsilon)(-2+N)(-1+N)N\pi(-1)^{2-k}}{2+\varepsilon} \times 2^{-2+\varepsilon} e^{-\frac{3\varepsilon\gamma}{2}} \eta^k \times$$

$$\frac{\Gamma(1-\frac{\varepsilon}{2}-i+j+k)\Gamma(-1-\frac{\varepsilon}{2})\Gamma(2+\frac{\varepsilon}{2})\Gamma(1+N)\Gamma(1+\varepsilon+i-k)\Gamma(-\frac{3\varepsilon}{2}+k)\Gamma(1-\varepsilon+k)\Gamma(3-\varepsilon+k)\Gamma(-\frac{1}{2}-\frac{\varepsilon}{2}+k)}{\Gamma(-\frac{3}{2}-\frac{\varepsilon}{2})\Gamma(\frac{5}{2}+\frac{\varepsilon}{2})\Gamma(2+i)\Gamma(1+k)\Gamma(2-i+j)\Gamma(2-\varepsilon+k)\Gamma(\frac{5}{2}-\varepsilon+k)\Gamma(-\frac{\varepsilon}{2}+k)\Gamma(5+\frac{\varepsilon}{2}+N)}$$

Example: a 2-mass 3-loop Feynman integral [arXiv:1804.02226]  
 (arose in the calculation of the gluonic operator matrix element  $A_{gg,Q}^{(3)}$ )



Mellin-Barnes-  
 and  $pF_q$ -technologies  $\rightarrow$

expression (95 MB) with

- 150 single sums
- 1000 double sums
- 12160 triple sums
- 1555 quadruple sums

Typical triple sum:

$$\sum_{j=0}^N \sum_{i=0}^j \sum_{k=0}^i \frac{(4+\varepsilon)(-2+N)(-1+N)N\pi(-1)^{2-k}}{2+\varepsilon} \times 2^{-2+\varepsilon} e^{-\frac{3\varepsilon\gamma}{2}} \eta^k \times$$


$$\frac{\Gamma(1-\frac{\varepsilon}{2}-i+j+k)\Gamma(-1-\frac{\varepsilon}{2})\Gamma(2+\frac{\varepsilon}{2})\Gamma(1+N)\Gamma(1+\varepsilon+i-k)\Gamma(-\frac{3\varepsilon}{2}+k)\Gamma(1-\varepsilon+k)\Gamma(3-\varepsilon+k)\Gamma(-\frac{1}{2}-\frac{\varepsilon}{2}+k)}{\Gamma(-\frac{3}{2}-\frac{\varepsilon}{2})\Gamma(\frac{5}{2}+\frac{\varepsilon}{2})\Gamma(2+i)\Gamma(1+k)\Gamma(2-i+j)\Gamma(2-\varepsilon+k)\Gamma(\frac{5}{2}-\varepsilon+k)\Gamma(-\frac{\varepsilon}{2}+k)\Gamma(5+\frac{\varepsilon}{2}+N)}$$

6 hours for this sum

$\sim$  10 years of calculation time for full expression

# Example: a 2-mass 3-loop Feynman integral [arXiv:1804.02226]

(arose in the calculation of the gluonic operator matrix element  $A_{gg,Q}^{(3)}$ )



Mellin-Barnes-  
and  ${}_pF_q$ -technologies  $\rightarrow$

expression (95 MB) with


- 150 single sums
- 1000 double sums
- 12160 triple sums
- 1555 quadruple sums

↓ SumProduction.m (2 hours)

expression (377 MB)  
consisting of 8 multi-sums

# Example: a 2-mass 3-loop Feynman integral [arXiv:1804.02226]

(arose in the calculation of the gluonic operator matrix element  $A_{gg,Q}^{(3)}$ )



Mellin-Barnes-  
and  $pF_q$ -technologies

expression (95 MB) with

- 150 single sums
- 1000 double sums
- 12160 triple sums
- 1555 quadruple sums

↓ SumProduction.m (2 hours)

expression (377 MB)  
consisting of 8 multi-sums


↓ EvaluateMultiSums.m

Example: a 2-mass 3-loop Feynman integral [arXiv:1804.02226]  
 (arose in the calculation of the gluonic operator matrix element  $A_{gg,Q}^{(3)}$ )

sum	size of sum (with $\varepsilon$ )	summand size of constant term	time of calculation	number of indef. sums
$\sum_{i_4=2}^{N-3} \sum_{i_3=0}^{i_4-2} \sum_{i_2=0}^{i_3} \sum_{i_1=0}^{\infty}$	17.7 MB	266.3 MB	177529 s (2.1 days)	1188
$\sum_{i_3=3}^{N-4} \sum_{i_2=0}^{i_3-1} \sum_{i_1=0}^{\infty}$	232 MB	1646.4 MB	980756 s (11.4 days)	747
$\sum_{i_2=3}^{N-4} \sum_{i_1=0}^{\infty}$	67.7 MB	458 MB	524485 s (6.1 days)	557
$\sum_{i_1=0}^{\infty}$	38.2 MB	90.5 MB	689100 s (8.0 days)	44
$\sum_{i_4=2}^{N-3} \sum_{i_3=0}^{i_4-2} \sum_{i_2=0}^{i_3} \sum_{i_1=0}^{i_2}$	1.3 MB	6.5 MB	305718 s (3.5 days)	1933
$\sum_{i_3=3}^{N-4} \sum_{i_2=0}^{i_3-1} \sum_{i_1=0}^{i_2}$	11.6 MB	32.4 MB	710576 s (8.2 days)	621
$\sum_{i_2=3}^{N-4} \sum_{i_1=0}^{i_2}$	4.5 MB	5.5 MB	435640 s (5.0 days)	536
$\sum_{i_1=3}^{N-4}$	0.7 MB	1.3 MB	9017s (2.5 hours)	68

# Example: a 2-mass 3-loop Feynman integral [arXiv:1804.02226]

(arose in the calculation of the gluonic operator matrix element  $A_{gg,Q}^{(3)}$ )



Mellin-Barnes-  
and  ${}_pF_q$ -technologies  $\rightarrow$

expression (95 MB) with

- 150 single sums
- 1000 double sums
- 12160 triple sums
- 1555 quadruple sums

↓ SumProduction.m (2 hours)

expression (377 MB)  
consisting of 8 multi-sums

↓ EvaluateMultiSums.m  
(3 month)

expression (154 MB)  
consisting of 4110 indefinite sums

**Example: a 2-mass 3-loop Feynman integral** [arXiv:1804.02226]  
 (arose in the calculation of the gluonic operator matrix element  $A_{gg,Q}^{(3)}$ )

Most complicated objects: generalized binomial sums, like


$$\sum_{h=1}^N 2^{-2h} (1-\eta)^h \binom{2h}{h} \left( \sum_{i=1}^h \frac{2^{2i} (1-\eta)^{-i}}{i \binom{2i}{i}} \right) \left( \sum_{i=1}^h \frac{(1-\eta)^i \binom{2i}{i}}{2^{2i}} \right) \times$$

$$\times \left( \sum_{i=1}^h \frac{2^{2i} (1-\eta)^{-i} \sum_{j=1}^i \frac{\sum_{k=1}^j (1-\eta)^k}{k}}{i \binom{2i}{i}} \right).$$



# Example: a 2-mass 3-loop Feynman integral [arXiv:1804.02226]

(arose in the calculation of the gluonic operator matrix element  $A_{gg,Q}^{(3)}$ )



Mellin-Barnes-  
and  $pF_q$ -technologies

expression (95 MB) with

- 150 single sums
- 1000 double sums
- 12160 triple sums
- 1555 quadruple sums

↓ SumProduction.m (2 hours)

expression (377 MB)  
consisting of 8 multi-sums


↓ EvaluateMultiSums.m  
(3 month)

expression (8.3 MB)  
consisting of  
74 indefinite sums

← Sigma.m (32 days)

expression (154 MB)  
consisting of 4110 indefinite sums

# Evaluation of Feynman Integrals



Behavior of particles



$$\int \Phi(N, \epsilon, x) dx$$

Feynman integrals

**DESY**  
(J. Blümlein)



$$\sum f(N, \epsilon, k)$$


complicated  
multi-sums

expression in  
special functions



**RISC**  
(Sigma-package)

# Evaluation of Feynman Integrals



Behavior of particles



$$\int \Phi(N, \epsilon, x) dx$$

Feynman integrals



LHC at CERN

**DESY**  
(J. Blümlein)



$$\sum f(N, \epsilon, k)$$

complicated  
multi-sums

applicable

expression in  
special functions

**RISC**

(Sigma-package)

